arXiv:0804.1990v2 [math.RT] 12 Jun 2009

Stein–Sahi complementary series and their degenerations

Yuri A. Neretin[1]

The paper is an introduction to the Stein–Sahi complementary series, and the unipotent representations. We also discuss some open problems related to these objects. For the sake of simplicity, we consider only the groups $\mathrm{U}(n,n)$.

1 Introduction

This paper[2] is an attempt to present an introduction to the Stein-Sahi complementary series available for non-experts and beginners.

1.1. History of the subject. Theory of infinite dimensional representations of semi-simple groups was initiated in pioneer works of I. M. Gelfand and M. A. Naimark (1946–1950), V. Bargmann [2] (1947), and K. O. Friedrichs [12] (1951–1953). The book [14] by I. M. Gelfand and M. A. Naimark (1950) contains a well-developed theory for complex classical groups $\mathrm{GL}(n,\mathbb{C})$, $\mathrm{SO}(n,\mathbb{C})$, $\mathrm{Sp}(2n,\mathbb{C})$ (the parabolic induction, complementary series, spherical functions, characters, Plancherel theorems). However, this classical book[3] contained various statements and asseverations that were not actually proved. In the modern terminology, some of chapters were 'mathematical physics'. The most of these statements were really proved by 1958–1962 in works of different authors (Harish-Chandra, F. A. Berezin, etc.).

In particular, I. M. Gelfand and M. A. Naimark (1950) claimed that they classified all unitary representations of $\mathrm{GL}(n,\mathbb{C})$, $\mathrm{SO}(n,\mathbb{C})$, $\mathrm{Sp}(2n,\mathbb{C})$. E. Stein [46] compared Gelfand–Naimark constructions for groups $\mathrm{SL}(4,\mathbb{C})\simeq\mathrm{SO}(6,\mathbb{C})$ and observed that they are not equivalent. In 1967 E. Stein constructed 'new' unitary representations of $\mathrm{SL}(2n,\mathbb{C})$.

D. Vogan [48] in 1986 obtained the classification of unitary representations of groups $\mathrm{GL}(2n)$ over real numbers $\mathbb{R}$ and quaternions $\mathbb{H}$. In particular, this work contains extension of Stein's construction to these groups. In 1990s, the Stein-type representations were a topic of interest of S. Sahi see [40], [41], [42], S. Sahi–E. Stein [44], A. Dvorsky–S. Sahi [8]–[9]. In particular, Sahi extended the construction to other series of classical groups, precisely to the groups $\mathrm{SO}(2n,2n)$, $\mathrm{U}(n,n)$, $\mathrm{Sp}(n,n)$, $\mathrm{Sp}(2n,\mathbb{R})$, $\mathrm{SO}^*(4n)$, $\mathrm{Sp}(4n,\mathbb{C})$, and $\mathrm{SO}(2n,\mathbb{C})$.

1.2. Stein–Sahi representations for $\mathrm{U}(n,n)$. Denote by $\mathrm{U}(n)$ the group of unitary $n\times n$-matrices. Consider the *pseudo-unitary* group $\mathrm{U}(n,n)$. We realize it as the group of $(n+n)\times(n+n)$-matrices $g=\begin{pmatrix}a&b\\c&d\end{pmatrix}$ satisfying the

---

[1] Supported by the grant FWF, project P19064, Russian Federal Agency for Nuclear Energy, Dutch grant NWO.047.017.015, and grant JSPS-RFBR-07.01.91209

[2] It is a strongly revised version of two sections of my preprint [30].

[3] Unfortunately the book exists only in Russian and German.

condition

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix}^{*} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

Lemma 1.1 *The formula*

$$z \mapsto z^{[g]} := (a+zc)^{-1}(b+zd) \tag{1.1}$$

determines an action of the group $\mathrm{U}(n,n)$ *on the space* $\mathrm{U}(n)$.

The unitary group is equipped by the Haar measure $d\mu(z)$, hence we can determine the Jacobian of a transformation (1.1) by

$$J(g,z) = \frac{d\mu(z^{[g]})}{d\mu(z)}.$$

Lemma 1.2 *The Jacobian of the transformation* $z \mapsto z^{[g]}$ *on* $\mathrm{U}(n)$ *is given by*

$$J(g,z) = |\det(a+zc)|^{-2n}.$$

Fix σ, $\tau \in \mathbb{C}$. For $g \in \mathrm{U}(n,n)$ we define the following linear operator in the space $C^\infty(\mathrm{U}(n))$:

$$\rho_{\sigma|\tau}(g)f(z) = f(z^{[g]})\det(a+zc)^{-n-\tau}\det\overline{(a+zc)}^{-n-\sigma}. \tag{1.2}$$

The formula includes powers of complex numbers, precise definition is given below. In fact, $g \mapsto \rho_{\sigma|\tau}(g)$ is a well-defined operator-valued function on the universal covering group $\mathrm{U}(n,n)^\sim$ of $\mathrm{U}(n,n)$.

The chain rule for Jacobians,

$$J(g_1g_2,z) = J(g_1,z)J(g_2,z^{[g_1]}), \tag{1.3}$$

implies

$$\rho_{\sigma|\tau}(g_1)\rho_{\sigma|\tau}(g_2) = \rho_{\sigma|\tau}(g_1g_2).$$

In other words, $\rho_{\sigma|\tau}$ is a linear representation of the group $\mathrm{U}(n,n)^\sim$.

Observation 1.3 *If* $\mathrm{Re}\,\sigma + \mathrm{Re}\,\tau = -n$, $\mathrm{Im}\,\sigma = \mathrm{Im}\,\tau$ *then a representation* $\rho_{\sigma|\tau}$ *is unitary in* $L^2(\mathrm{U}(n))$.

This easily follows from the formula for the Jacobian.

Next, let σ, τ be real. We define the Hermitian form on $C^\infty(\mathrm{U}(n))$ by the formula

$$\langle f_1, f_2\rangle_{\sigma|\tau} := \int_{\mathrm{U}(n)}\int_{\mathrm{U}(n)} \det(1-zu^*)^\sigma(1-z^*u)^\tau f_1(z)\,\overline{f_2(u)}\,d\mu(z)\,d\mu(u). \tag{1.4}$$

Proposition 1.4 *The operators* $\rho_{\sigma|\tau}(g)$ *preserve the Hermitian form* $\langle\cdot,\cdot\rangle_{\sigma|\tau}$.

Theorem 1.5 *For σ, $\tau \notin \mathbb{Z}$, the Hermitian form $\langle \cdot, \cdot \rangle_{\sigma|\tau}$ is positive iff integer parts of numbers $-\sigma - n$ and τ are equal.*

In fact, the domain of positivity is the square $-1 < \tau < 0$, $-n < \sigma < -n+1$ and its shifts by vectors $(-j, j)$, $j \in \mathbb{Z}$, see Figure 5.

In particular, under this condition, a representation $\rho_{\sigma|\tau}$ is unitary.

For some values of (σ, τ) the form $\langle \cdot, \cdot \rangle_{\sigma|\tau}$ is positive semi-definite. There are two the most important cases.

1. For $\tau = 0$, we get highest weight representations (or holomorphic representations). Thus, the Stein–Sahi representations are nearest relatives of holomorphic representations.

2. For $\tau = 0$, $\sigma = 0, -1, -2, \dots, -n$ we obtain some exotic 'small' representations of $\mathrm{U}(n, n)$.

1.3. The structure of the paper. *We discuss only groups*[4] $\mathrm{U}(n, n)$.

In Section 2 we consider the case $n = 1$ and present the Pukanszky classification [37] of unitary representations of the universal covering group of $\mathrm{SL}(2, \mathbb{R}) \simeq \mathrm{SU}(1, 1)$.

In Section 3 we discuss Stein-Sahi representations of arbitrary $\mathrm{U}(n, n)$. In Section 4 we explain relations of Stein–Sahi representations and holomorphic representations. In Section 5 we give explicit constructions of the Sahi 'unipotent' representations.

In Section 6 we discuss some open problems of harmonic analysis.

1.4. Notation. Let a, u, $v \in \mathbb{C}$. Denote

$$a^{\{u|v\}} := a^u \overline{a}^v. \qquad (1.5)$$

If $u - v \in \mathbb{Z}$, then this expression is well defined for all $a \neq 0$. However, the expression is well defined in many other situations, for instance if $|1 - a| < 1$ and u, v are arbitrary (and even for $|1 - a| = 1$, $a \neq 1$)

The *norm* $\|z\|$ of an $n \times n$-matrix z is the usual norm of a linear operator in the standard Euclidean space $\mathbb{C}^n$.

---

[4] A comment for experts. Stein–Sahi representations of a semisimple Lie group G are complementary series induced from a maximal parabolic subgroup with Abelian nilpotent radical.

The cases $G = \mathrm{U}(n, n)$, $\mathrm{Sp}(2n, \mathbb{R})$, $G = \mathrm{SO}^*(4n)$ (related to tube type Hermitian symmetric spaces) are parallel. The only difficulty is Theorem 3.11 (the expansion of the integral kernel in characters, we choose $G = \mathrm{U}(n, n)$, because this can be done by elementary tools). In the general Hermitian case, one can refer to the version of the Kadell integral [20] from [29] (the integrand is a product of a Jack polynomial and a Selberg-type factor.

For other series of groups, Stein-Sahi representations depend on one parameter, and picture is more pure (in particular, inner products for degenerate ('unipotent') representations can be written immediately). A BC-analog of Kadell integral is unknown (certainly, it must exist, and some special cases were evaluated in the literature, see e.g.,[30]). On the other hand, Stein-Sahi representations have multiplicity free K-spectra. In such situation, there is lot of ways for examination of positivity of inner products, see e.g. [41], [42], [5].

New elements of this paper are 'blow-up construction' for unipotent representations and (apparently) tame models for representations of universal coverings. The representations themselves were constructed in works of Sahi.

We denote the Haar measure on the unitary group $\mathrm{U}(n)$ by μ; assume that the complete measure of the group is 1.

The *Pochhammer symbol* is given by

$$(a)_n := \frac{\Gamma(a+n)}{\Gamma(a)} = \begin{cases} a(a+1)\dots(a+n-1) & \text{if } n \geqslant 0 \\ \frac{1}{(a-1)\dots(a-n)} & \text{if } n < 0. \end{cases} \tag{1.6}$$

2 Unitary representations of $\mathrm{SU}(1,1)$

Denote by $\mathrm{SU}(1,1)^\sim$ the universal covering group of $\mathrm{SU}(1,1)$.

In this section, we present constructions of all irreducible unitary representations of $\mathrm{SU}(1,1)^\sim$. According the Bargmann–Pukanszky theorem there are 4 types of such representations:

a) unitary principal series;

b) complementary series;

c) highest weight and lowest weight representations;

d) The one-dimensional representation.

Models of these representations are given below.

The general Stein–Sahi representations are a strange 'higher copy' of the $\mathrm{SU}(1,1)$-picture.

References. The classification of unitary representations of $\mathrm{SL}(2,\mathbb{R}) \simeq \mathrm{SU}(1,1)$ was obtained by V. Bargmann [2]; it was extended to the $\mathrm{SU}(1,1)^\sim$ by L. Pukanszky [37], see also P. Sally [45]. □

A. Preliminaries

2.1. Fourier series and distributions. By S^1 we denote the unit circle $|z| = 1$ in the complex plane $\mathbb{C}$. We parameterize S^1 by $z = e^{i\varphi}$.

By $C^\infty(S^1)$ we denote the space of smooth functions on S^1. Recall, that

$$f(\varphi) = \sum_{n=-\infty}^{\infty} a_n e^{in\varphi} \in C^\infty(S^1) \qquad \text{iff } |a_n| = o(|n|^{-L}) \text{ for all } L.$$

Recall that a distribution $h(\varphi)$ on the circle admits an expansion into a Fourier series,

$$h(\varphi) = \sum_{n=0}^{\infty} b_n e^{in\varphi}, \quad \text{where } |b_n| = O(|n|^L) \text{ for some } L.$$

For $s \in \mathbb{R}$ we define the *Sobolev space* $W^s(S^1)$ as the space of distributions

$$h(\varphi) = \sum_{n=0}^{\infty} b_n e^{in\varphi} \quad \text{such that } \sum |b_n|^2 (1+|n|)^{2s} < \infty.$$

By definition, $W^0(S^1) = L^2(S^1)$. For positive integer $s = k$ this condition is equivalent $\frac{\partial^k}{\partial\varphi^k} h \in L^2(S^1)$. Evidently, $s < s'$ implies $W^s \supset W^{s'}$.

2.2. The group $\mathrm{SU}(1,1)$**.** The group $\mathrm{SU}(1,1) \simeq \mathrm{SL}(2,\mathbb{R})$ consists of all complex 2×2-matrices having the form

$$g = \begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix}, \qquad \text{where } |a|^2 - |b|^2 = 1.$$

This group acts on the disc $|z| < 1$ and on the circle $|z| = 1$ by the *Möbius transformations*

$$z \mapsto (a + \overline{b}z)^{-1}(b + \overline{a}z).$$

2.3. A model of the universal covering group $\mathrm{SU}(1,1)^\sim$**.** Recall that the fundamental group of $\mathrm{SU}(1,1)$ is $\mathbb{Z}$. A loop generating the fundamental group is

$$\mathfrak{R}(\varphi) = \begin{pmatrix} e^{i\varphi} & 0 \\ 0 & e^{-i\varphi} \end{pmatrix}, \qquad \mathfrak{R}(2\pi) = \mathfrak{R}(0) = 1. \tag{2.1}$$

Some example of multi-valued continuous function on $\mathrm{SU}(1,1)$ are

$$\begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix} \mapsto \ln a, \qquad \begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix} \mapsto a^\lambda := a^{\lambda \ln a}.$$

We can realize $\mathrm{SU}(1,1)^\sim$ as a subset in $\mathrm{SU}(1,1) \times \mathbb{C}$ consisting of pairs

$$\left(\begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix}, \sigma \right), \quad \text{where } e^\sigma = a.$$

Thus, for a given matrix $\begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix}$ the parameter σ ranges if the countable set $\sigma = \ln a + 2\pi k i$.

Define a multiplication in $\mathrm{SU}(1,1) \times \mathbb{C}$ by

$$(g_1, \sigma_1) \circ (g_2, \sigma_2) = (g_1 g_2, \sigma_1 + \sigma_2 + c(g_1, g_2)),$$

where $c(g_1, g_2)$ is the *Berezin–Guichardet* cocycle,

$$c(g_1, g_2) = \ln \frac{a_3}{a_1 a_2},$$

Here a_3 is the matrix element of $g_3 = g_1 g_2$.

Theorem 2.1 *a)* $\left| \frac{a_3}{a_1 a_2} - 1 \right| < 1$*, and therefore the logarithm is well defined.*

b) The operation $\circ$ *determines the structure of a group on* $\mathrm{SU}(1,1) \times \mathbb{C}$*.*

c) $\mathrm{SU}(1,1)^\sim$ *is a subgroup in the latter group.*

The proof is a simple and nice exercise.

Now we can define the single-valued function $\ln a$ on $\mathrm{SU}(1,1)$ by setting $\ln a := \sigma$.

B. Non-unitary and unitary principal series

2.4. Principal series of representations of $\mathrm{SU}(1,1)$. Fix p, $q \in \mathbb{C}$. For $g \in \mathrm{SU}(1,1)$ define the operator $T_{p|q}(g)$ in the space $C^\infty(S^1)$ by the formula

$$T_{p|q}\begin{pmatrix} a & b \\ \overline{b} & \overline{a} \end{pmatrix} f(z) = f\Bigl(\frac{b+\overline{a}z}{a+\overline{b}z}\Bigr)(a+\overline{b}z)^{\{-p|-q\}}, \tag{2.2}$$

here we use the notation (1.5) for complex powers.

Observation 2.2 *a) $T_{p|q}$ is a well-defined operator-valued function on* $\mathrm{SU}(1,1)^\sim$.

b) It satisfies

$$T_{p|q}(g_1)T_{p|q}(g_2) = T_{p|q}(g_1 g_2).$$

Proof. a) First,

$$(a+\overline{b}z)^{-p}\,\overline{(a+\overline{b}z)}^{\;-q} = a^{-p}\cdot\overline{a}^{\;-q}(1+a^{-1}\overline{b}z)^{-p}\,\overline{(1+a^{-1}\overline{b}z)}^{\;-q}.$$

Since $|z|=1$ and $|a|>|b|$, the last two factors are well defined. Next,

$$a^{-p}\,\overline{a}^{\;-q} := \exp\Bigl\{-p\ln a + q\,\overline{\ln a}\Bigr\}$$

and $\ln a$ is a well-defined function on $\mathrm{SU}(1,1)^\sim$.

Proof of b). One can verify this identity for g_1, g_2 near the unit and refer to the analytic continuation. □

The representations $T_{p|q}(g)$ are called *representations of the principal (non-unitary) series.*

Remark. a) A representation $T_{p|q}$ is a single-valued representation of $\mathrm{SU}(1,1)$ iff $p-q$ is integer.

2.5. The action of the Lie algebra. The Lie algebra $\mathfrak{su}(1,1)$ of $\mathrm{SU}(1,1)$ consists of matrices

$$\begin{pmatrix} i\alpha & \beta \\ \overline{\beta} & -i\alpha \end{pmatrix}, \qquad \text{where } \alpha\in\mathbb{R},\ \beta\in\mathbb{C}.$$

It is convenient to take the following basis in the complexification $\mathfrak{su}(1,1)_\mathbb{C} = \mathfrak{sl}(2,\mathbb{C})$:

$$L_0 := \frac{1}{2}\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad L_- := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad L_+ := \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} \tag{2.3}$$

These generators act in $C^\infty(S^1)$ by the following operators

$$L_0 = z\frac{d}{dz} + \frac{1}{2}(p-q), \qquad L_- = \frac{d}{dz} - qz^{-1}, \qquad L_+ = z^2\frac{d}{dz} + pz. \tag{2.4}$$

Equivalently,

$$L_0 z^n = \big(n + \frac{1}{2}(p-q)\big) z^n, \qquad L_- z^n = (n-q) z^{n-1}, \qquad L_+ z^n = (n+p) z^{n+1}. \tag{2.5}$$

2.6. Subrepresentations.

Proposition 2.3 *A representation $T_{p|q}$ is irreducible iff $p,\ q \notin \mathbb{Z}$.*

PROOF. Let $p,\ q \notin \mathbb{Z}$. Consider an L_0-eigenvector z^n. Then all vectors $(L_+)^k z^n$, $(L_-)^l z^n$ are nonzero. They span the whole space $C^\infty(S^1)$. ☐

Observation 2.4 *a) If $q \in \mathbb{Z}$, then z^q, z^{q+1}, ... span a subrepresentation in $T_{p|q}$.*

b) If $p \in \mathbb{Z}$, then z^{-p}, z^{-p-1}, z^{-p-2}, ... span a subrepresentation in $T_{p|q}$.

PROOF OF A). Clearly, our subspace is L^0-invariant and L^+-invariant. On the other hand, $L^- z^q = 0$, and we can not leave our subspace. ☐

All possible positions of subrepresentations of $T_{p|q}$ are listed on Figure 1.

2.7. Shifts of parameters.

Observation 2.5 *If k is integer, then $T_{p+k|q-k} \simeq T_{p|q}$. The intertwining operator is*

$$Af(z) = z^k f(z).$$

A verification is trivial. ☐

2.8. Duality. Consider the bilinear map

$$\Pi : C^\infty(S^1) \times C^\infty(S^1) \to \mathbb{C}$$

given by

$$(f_1, f_2) \mapsto \frac{1}{2\pi} \int_0^{2\pi} f_1(e^{i\varphi}) f_2(e^{i\varphi})\, d\varphi = \frac{1}{2\pi} \int_0^{2\pi} f_1(z) f_2(z)\, \frac{dz}{z}. \tag{2.6}$$

Observation 2.6 *Representations $T_{p|q}$ and $T_{1-p|1-q}$ are dual with respect to Π, i.e.,*

$$\Pi\left(T_{p|q}(g) f_1, T_{1-p|1-q}(g) f_2\right) = \Pi(f_1, f_2). \tag{2.7}$$

PROOF. After simple cancelations we get the following expression in the left hand side of (2.7)

$$\frac{1}{2\pi i} \int_{|z|=1} f_1\left(\frac{b+\overline{a}z}{a+\overline{b}z}\right) f_2\left(\frac{b+\overline{a}z}{a+\overline{b}z}\right) \cdot (a+\overline{b}z)^{-1} \overline{(a+\overline{b}z)}^{\,-1} \frac{dz}{z}.$$

Keeping in mind $\overline{z} = z^{-1}$, we transform

$$(a+\overline{b}z)^{-1} \overline{(a+\overline{b}z)}^{\,-1} \frac{dz}{z} = (a+\overline{b}z)^{-1} (\overline{b}+az)^{-1}\, dz = \left(\frac{b+\overline{a}z}{a+\overline{b}z}\right)^{-1} d\left(\frac{b+\overline{a}z}{a+\overline{b}z}\right).$$



a) *q is integer*;



b) *p is integer*;



c) *p, q are integer*, $q + p \geqslant 1$;



d) *p, q are integer*, $q + p \leqslant 1$.

Figure 1: Subrepresentations of the principal series. Black circles enumerate vectors z^n. A representation $T_{p|q}$ is reducible iff $p \in \mathbb{Z}$ or $q \in \mathbb{Z}$.

Now the integral comes to the desired form:

$$\frac{1}{2\pi i}\int_{|u|=1} f_1(u)\, f_2(u)\,\frac{du}{u}. \qquad \square$$

We also define a sesquilinear map

$$\Pi^* : C^\infty(S^1)\times C^\infty(S^1)\to \mathbb{C}$$

by

$$\Pi^*(f_1,f_2) := \Pi(f_1,\overline{f}_2) = \int_0^{2\pi} f_1(z)\overline{f_2(z)}\,\frac{dz}{z}. \tag{2.8}$$

Observation 2.7 *Representations* $T_{p|q}$ *and* $T_{1-\overline{q}|1-\overline{p}}$ *are dual with respect to* Π^**.*

Proof is same. $\square$

2.9. Intertwining operators. Consider the integral operator

$$I_{p|q} : C^\infty(S^1)\to C^\infty(S^1)$$

given by

$$I_{p|q}f(u) = \frac{1}{2\pi i\,\Gamma(p+q-1)}\int_{|z|=1}(1-z\overline{u})^{\{p-1|q-1\}}f(z)\,\frac{dz}{z}, \tag{2.9}$$

where the function $(1-z\overline{u})^{\{p-1|q-1\}}$ is defined by

$$(1-z\overline{u})^{\{p-1|q-1\}} := \lim_{t\to 1^-}(1-tz\overline{u})^{\{p-1|q-1\}} \tag{2.10}$$

The integral converges if $\mathrm{Re}(p+q) > -1$.

Theorem 2.8 *The map* $(p|q)\mapsto I_{p|q}$ *admits the analytic continuation to a holomorphic operator-valued function on* $\mathbb{C}^2$*.*

Theorem 2.9 *The operator* $I_{p|q}$ *intertwines* $T_{p|q}$ *and* $T_{1-q|1-p}$*, i.e.,*

$$T_{1-p|1-q}(g)\, I_{p|q} = I_{p|q}\, T_{p|q}(g).$$

Corollary 2.10 *If* $p\notin\mathbb{Z}$*,* $q\notin\mathbb{Z}$*, then the representations* $T_{p|q}$ *and* $T_{1-q|1-p}$ *are equivalent.*

2.10. Proof of Theorems 2.8, 2.9.

Lemma 2.11 *The expansion of the distribution (2.10) into the Fourier series is given by*

$$(1-z\overline{u})^{p-1}(1-\overline{z}u)^{q-1} = \frac{\Gamma(p+q-1)}{\Gamma(p)\Gamma(q)}\sum_{n=-\infty}^{\infty}\frac{(1-q)_n}{(p)_n}\left(\frac{z}{u}\right)^n = \tag{2.11}$$

$$= \Gamma(p+q-1)\sum_{n=-\infty}^{\infty}\frac{(-1)^n}{\Gamma(p+n)\Gamma(q-n)}\left(\frac{z}{u}\right)^n. \tag{2.12}$$

PROOF. Let $\operatorname{Re} p$, $\operatorname{Re} q$ be sufficiently large. Then we write

$$(1-z\overline{u})^{p-1}(1-\overline{z}u)^{q-1} = \Bigl[\sum_{j\geqslant 0}\frac{(1-p)_j}{j!}\Bigl(\frac{z}{u}\Bigr)^j\Bigr]\cdot\Bigl[\sum_{l\geqslant 0}\frac{(1-q)_l}{l!}\Bigl(\frac{u}{z}\Bigr)^l\Bigr] \qquad (2.13)$$

and open brackets in (2.13). For instance, the coefficient at $(z/u)^0$ is

$$\sum_{k\geqslant 0}\frac{(1-p)_k(1-q)_k}{k!\,k!} = {}_2F_1(1-p,1-q;1;1),$$

where ${}_2F_1$ is the Gauss hypergeometric function. We evaluate the sum with the Gauss summation formula for ${}_2F_1(1)$, see [18], (2.1.14). □

PROOF OF THEOREM 2.8. Denote by

$$c_n := \frac{(-1)^n}{\Gamma(p+n)\Gamma(q-n)}$$

the Fourier coefficients in (2.12). Evidently, c_n admits holomorphic continuation to the whole plane[5] $\mathbb{C}^2$.

By [18], (1.18.4),

$$\frac{\Gamma(n+a)}{\Gamma(n+b)} \sim |n|^{a-b} \qquad \text{as } n\to\pm\infty.$$

Keeping in mind (2.11), we get

$$c_n \sim \text{const}\cdot|n|^{1-p-q} \qquad \text{as } n\to\pm\infty. \qquad (2.14)$$

Then

$$I_{p|q} : z^n \mapsto c_{-n}z^n$$

and

$$I_{p|q} : \sum a_n z^n \mapsto \sum a_n c_{-n} z^n.$$

Obviously, this map sends smooth functions to smooth functions. □

PROOF OF COROLLARY 2.10. In this case, all $c_n \neq 0$. □

PROOF OF THEOREM 2.9. The calculation is straightforward,

$$T_{1-q|1-p}(g)I_{p|q}f(u) = \\ = \frac{1}{2\pi i}(a+\overline{b}u)^{\{q-1|p-1\}}\int_{|u|=1}\Bigl(1-\Bigl(\frac{b+\overline{a}u}{a+\overline{b}u}\Bigr)\overline{z}\Bigr)^{\{q-1|p-1\}} f(z)\,\frac{dz}{z}.$$

Next, we observe

$$(a+\overline{b}u)\Bigl(1-\Bigl(\frac{b+\overline{a}u}{a+\overline{b}u}\Bigr)\overline{z}\Bigr) = (a-b\overline{z})\Bigl(1-u\Bigl(\frac{-\overline{b}+\overline{az}}{a-b\overline{z}}\Bigr)\Bigr)$$

---

[5] The Gamma function $\Gamma(z)$ has simple poles at $z=0,-1,-2,\dots$ and does not have zeros. Therefore $1/(\Gamma(p+n)\Gamma(q-n))$ has zeros at $p=-n,-n-1,\dots$ and at $q=n,n-1,\dots$.

In particular, if both p, q are integer and $q<p$, when $I_{p|q}=0$.



Figure 2: The unitary principal series in coordinates
$h=(p-q+1)/2$, $s=\frac{1}{i}(p+q-1)/2$.
Equivalently,
$p=h+is$, $q=1-h+is$.
The shift $h\mapsto h+1$ does not change a representation. Also the symmetry $s\mapsto -s$ sends a representation to an equivalent one. Therefore representations of the principal series are enumerated by the a semi-strip $0\leqslant h<1$, $s\geqslant 0$. It is more reasonable to think that representations of the unitary principal series are enumerated by points of a semi-cylinder (s,h), where $s\geqslant 0$ and h is defined modulo equivalence $h\sim h+k$, where $h\in\mathbb{Z}$.

and come to

$$\frac{1}{2\pi i}\int_{|z|=1}\left(1-u\left(\frac{-\overline{b}+\overline{a}\overline{z}}{a-b\overline{z}}\right)\right)^{\{q-1|p-1\}}(a-b\overline{z})^{\{q-1|p-1\}}f(z)\,\frac{dz}{z}.$$

Now we change a variable again

$$z=\frac{b+\overline{a}w}{a+\overline{b}w},\qquad \overline{w}=\frac{-\overline{b}+\overline{a}\,\overline{z}}{a-b\overline{z}}$$

and come to the desired expression

$$\frac{1}{2\pi i}\int_{|w|=1}(1-u\overline{w})^{\{p-1|q-1\}}f\left(\frac{b+\overline{a}w}{a+\overline{b}w}\right)(a+\overline{b}w)^{\{-p|-q\}}\,\frac{dw}{w}.$$

2.11. The unitary principal series.

Observation 2.12 *A representation $T_{p|q}$ is unitary in $L^2(S^1)$ iff*

$$\operatorname{Im}p=\operatorname{Im}q,\qquad \operatorname{Re}p+\operatorname{Re}q=1. \tag{2.15}$$

PROOF is straightforward, also this follows from Observation 2.7. □

C. The complementary series

2.12. The complementary series. Now let

$$0<p<1\qquad 0<q<1. \tag{2.16}$$

Consider the Hermitian form on $C^\infty(S^1)$ given by

$$\langle f_1, f_2\rangle_{p|q} = \frac{1}{(2\pi i)^2\,\Gamma(p+q-1)} \int\limits_{|z|=1} \int\limits_{|u|=1} (1 - z\overline{u})^{\{p-1|q-1\}} f_1(z)\overline{f_2(u)}\,\frac{dz}{z}\frac{du}{u}. \tag{2.17}$$

By (2.12),

$$\langle z^n, z^m\rangle_{p|q} = \frac{1}{\Gamma(p)\Gamma(q)}\frac{(1-q)_n}{(p)_n}\cdot\delta_{m,n}. \tag{2.18}$$

Theorem 2.13 *If $0 < p < 1$, $0 < q < 1$, then the inner product (2.17) is positive definite.*

Proof. Indeed, in this case all coefficients

$$\frac{(1-q)_n}{(p)_n} = \frac{(1-p)_{-n}}{(q)_{-n}}$$

in (2.17) are positive. □

Theorem 2.14 *Let $0 < p < 1$, $0 < q < 1$. Then the representation $T_{p|q}$ is unitary with respect to the inner product $\langle\cdot,\cdot\rangle_{p|q}$, i.e.,*

$$\langle T_{p|q}(g)f_1, T_{p|q}(g)f_2\rangle_{p|q} = \langle f_1, f_2\rangle_{p|q}.$$

Proof. This follows from Theorem 2.9 and Observation 2.7. Indeed,

$$\langle f_1, f_2\rangle_{p|q} = \Pi^*(I_{p|q}f_1, f_2)$$

and

$$\begin{aligned}\Pi^*(I_{p|q}T_{p|q}(g)f_1, T_{p|q}(g)f_2) = \Pi^*(T_{1-q|1-p}(g)I_{p|q}f_1, T_{p|q}(g)f_2) =\\ = \Pi^*(I_{p|q}f_1, f_2) = \langle f_1, f_2\rangle_{p|q}.\end{aligned}$$

Keeping in mind our future purposes, we propose another (homotopic) proof. Substitute

$$z = \frac{b+\overline{a}z'}{a+\overline{b}z'}, \qquad u = \frac{b+\overline{a}u'}{a+\overline{b}u'}$$

to the integral in (2.17). Applying the identity

$$1 - \left(\frac{b+\overline{a}z'}{a+\overline{b}z'}\right)\overline{\left(\frac{b+\overline{a}u'}{a+\overline{b}u'}\right)} = (a+\overline{b}z')^{-1}(1 - z'\overline{u}')\overline{(a+\overline{b}u')}^{-1},$$

we get

$$\langle T_{p|q}(g)f_1, T_{p|q}(g)f_2\rangle_{p|q}.$$ □

2.13. Sobolev spaces. Denote by $\mathcal{H}_{p|q}$ the completion of $C^\infty(S^1)$ with respect to the inner product of the complementary series.



Figure 3: The complementary series. The diagonal is contained in the principal series (the segment of the axis Oh on Fig.2). The symmetry with respect to the diagonal sends a representation to an equivalent representation.

First, we observe that the principal series and the complementary series have an intersection, see (2.15), (2.16), namely the interval

$$p + q = 1, \qquad 0 < p < 1.$$

In this case the inner product (2.18) is the L^2-inner product, i.e., $\mathcal{H}_{p|1-p} \simeq L^2(S^1)$.

Next consider arbitrary (p, q), where $0 < p < 1$, $0 < q < 1$. By (2.14), the space $\mathcal{H}_{p|q}$ consists of Fourier series $\sum a_n z^n$ such that

$$\sum_{n=-\infty}^{\infty} |a_n|^2 n^{1-p-q} < \infty.$$

Thus, $\mathcal{H}_{p,q}$ is the Sobolev space $W^{(1-p-q)/2}(S^1)$.

D. Holomorphic and anti-holomorphic representations

Denote by D the disk $|z| < 1$ in $\mathbb{C}$.

2.14. Holomorphic (highest weight) representations. Set $q = 0$,

$$T_{p|0} f(z) = f\left(\frac{b + \overline{a} z}{a + \overline{b} z}\right) (a + \overline{b} z)^{-p}.$$

Since $|a| > |b|$, the factor $(a + \overline{b} z)^{-p}$ is holomorphic in the disk D. Therefore the space of holomorphic functions in D is $\mathrm{SU}(1,1)^\sim$-invariant. Denote the representation of $\mathrm{SU}(1,1)^\sim$ in the space of holomorphic functions by T_p^+.

Theorem 2.15 *a) For $p > 0$ the representation T_p^+ is unitary, the invariant inner product in the space of holomorphic functions is*

$$\Big\langle \sum_{n \geqslant 0} a_n z^n, \sum_{n \geqslant 0} b_n z^n \Big\rangle = \sum_{n > 0} \frac{n!}{(p)_n} a_n \overline{b}_n. \tag{2.19}$$

b) For $p > 1$ the invariant inner product admits the following integral representation:

$$\langle f_1, f_2\rangle = \frac{p-1}{\pi}\iint_{|z|<1} f_1(z)\overline{f_2(z)}\,(1-|z|^2)^{p-2}d\lambda(z),$$

where $d\lambda(z)$ is the Lebesgue measure in the disk.

c) For $p = 1$ the invariant inner product is

$$\langle f_1, f_2\rangle = \frac{1}{2\pi}\int_0^{2\pi} f_1(e^{i\varphi})\overline{f_2(e^{i\varphi})}\,d\varphi = \frac{1}{2\pi i}\int_{|z|=1} f_1(z)\overline{f_2(z)}\,\frac{dz}{z}. \qquad (2.20)$$

We denote this Hilbert space of holomorphic functions by $\mathcal{H}_p^+$.

PROOF. The invariance of inner products in b), c) can be easily verified by straightforward calculations.

To prove a), we note that weight vectors z^n must be pairwise orthogonal.

Next, operators of the Lie algebra $\mathfrak{su}(1,1)$ must be skew-self-adjoint. The generators of the Lie algebra must satisfy

$$(L_+)^* = L_-.$$

Therefore,

$$\langle L_+z^n, z^{n+1}\rangle = \langle z^n, L_-z^{n+1}\rangle$$

or

$$(n+p)\langle z^{n+1}, z^{n+1}\rangle = (n+1)\langle z^n, z^n\rangle.$$

This implies a).

If $p = 1$, then $\langle z^n, z^n\rangle = 1$ for $n \geqslant 0$, i.e., we get the L^2-inner product. □

The theorem does not provide us an explicit integral formula for inner product in $\mathcal{H}_p^+$ if $0 < p < 1$. There is another way of description of inner products in spaces of holomorphic functions.

2.15. Reproducing kernels.

Theorem 2.16 *For each $p > 0$, for any $f \in \mathcal{H}_p^+$, and for each $a \in D$*

$$\langle f(z), (1 - z\overline{a})^{-p}\rangle = f(a) \qquad \textit{(the reproducing property).} \qquad (2.21)$$

PROOF. Indeed,

$$\langle\sum a_n z^n, \sum \frac{(p)_n}{n!} z^n\overline{u}^n\rangle = \sum a_n \frac{(p)_n}{n!} u^n \langle z^n, z^n\rangle = \sum a_n u^n = f(u). \qquad \square$$

In fact, the identity (2.21) is an all-sufficient definition of the inner product. We will not discuss this (see [10], [31]), and prefer another way.

2.16. Realizations of holomorphic representations in quotient spaces. Consider the representation $T_{-1|-1-p}$ of the principal series,

$$T_{-1|-1-q} f(z) = f\left(\frac{b+\overline{a}z}{a+\overline{b}z}\right)(a+\overline{b}z)^{-1}\overline{(a+\overline{b}z)}^{-1-p}.$$

The corresponding invariant Hermitian form in $C^\infty(S^1)$, is

$$\langle f_1, f_2\rangle_{-1|-1-p} = \frac{1}{(2\pi i)^2}\int_{|z|=1}\int_{|u|=1}(1-\overline{z}u)^{-p} f_1(z)\,\overline{f_2(u)}\,\frac{dz}{z}\,\frac{du}{u}. \tag{2.22}$$

(we write another pre-integral factor in comparison with (2.17)). The integral diverges for $p > 1$. However, we can define the inner product by

$$\langle z^n, z^n\rangle = \begin{cases} \frac{(p)_n}{n!} & \text{if } n \geqslant 0 \\ 0 & \text{if } n < 0 \end{cases},$$

the latter definition is valid for all $p > 0$.

We denote by $L \subset C^\infty(S^1)$ the subspace consisting of series $\sum_{n<0} a_n z^n$. This subspace is SU$(1,1)$-invariant and our form is nondegenerate and positive definite on the quotient space $C^\infty(S^1)/L$.

Next, we consider the intertwining operator

$$\widetilde{I}_{-1|-1-p} : C^\infty(S^1) \to C^\infty(S^1)$$

as above (but we change a normalization of the integral),

$$\widetilde{I}_{-1|-1-p} f(u) = \frac{1}{2\pi i}\int_{|z|=1}(1-\overline{z}u)^{-p} f(z)\,\frac{dz}{z}$$

The kernel of the operator is L and the image consists of holomorphic functions.

Observation 2.17 *a) The operator $I_{-1|-1-p}$ is a unitary operator*

$$C^\infty(S^1)/L \to \mathcal{H}_p^+.$$

b) The representation $T_{-1|-1-p}$ in $C^\infty(S^1)/L$ is equivalent to the highest weight representation T_p^+

2.17. Lowest weight representations. Now set $p = 0$, $q > 0$. Then operators $T_{0|q}$ preserve the subspace consisting of 'antiholomorphic' functions $\sum_{n\leqslant 0} a_n z^n$. Denote by T_q^- the corresponding representation in the space of antiholomorphic functions. These representations are unitary.

We omit further discussion because these representations are twins of highest weight representations.

E. The blow-up trick

Here we discuss a trick that produces 'unipotent' representations of U(n,n) for $n \geqslant 2$, see Subsection 5.2.

2.18. The exotic case $p = 1$, $q = 0$. In this case,

$$T_{1|0} = T_1^+ \oplus T_1^-.$$



Figure 4: a) The structure of the representation $T_{1|0}$.
b) Ways to $(p,q)=(1,0)$ from different directions give origins to different invariant Hermitian forms on $T_{1|0}$. By our normalization, the inner product is positive definite in the gray triangle and negative definite in the white triangle. Therefore coming to $(1,0)$ from the grey triangle we get a positive form.

Let us discuss the behavior of the inner product of the complementary series near the point $(p|q)=(1|0)$,

$$\langle f_1, f_2\rangle_{p|q} = \frac{1}{(2\pi i)^2}\int_{|z|=1}(1-z\overline{u})^{\{p-1|q-1\}} f_1(z)\overline{f_2(z)}\,\frac{dz}{z}. \tag{2.23}$$

Consider the limit of this expression as $p\to 1$, $q\to 0$. The Fourier coefficients of the kernel are the following meromorphic functions

$$c_n(p,q)=\frac{(-1)^n\Gamma(p+q-1)}{\Gamma(q-n)\Gamma(p+n)}.$$

Note that

1. $c_n(p,q)$ has a pole at the line $p+q=1$;
2. for $n\geqslant 0$, the function $c_n(p,q)$ has a zero on the line $q=0$;
3. for $n<0$, the function $c_n(p,q)$ has a zero at the line $p=0$.

Thus our point $(p,q)=(1,0)$ lies on the intersection of a pole and of a zero of the function $c_n(p,q)$. Let us substitute

$$p=1+\varepsilon s \qquad q=\varepsilon t, \qquad \text{where } s+t\neq 0$$

to $c_n(p,q)$ and pass to the limit as $\varepsilon\to 0$. Recall that

$$\Gamma(z)=\frac{(-1)^n}{n!\,(z+n)}+O(1), \qquad \text{as } z\to -n, \text{ where } n=0,\,1,\,2,\ldots \tag{2.24}$$

Therefore we get

$$\lim_{\varepsilon\to 0} c_n(1+\varepsilon s, \varepsilon t) = \begin{cases} \frac{t}{t+s} & \text{if } n \geqslant 0 \\ -\frac{s}{t+s} & \text{if } n < 0. \end{cases}$$

In particular, for $s = 0$ we get T_1^+-inner product, and for $t = 0$ we get T_1^--inner product. Generally,

$$\lim_{\varepsilon\to 0} \langle \sum_{n=-\infty}^{\infty} a_n z^n, \sum_{n=-\infty}^{\infty} b_n z^n \rangle_{1+\varepsilon s|\varepsilon t} = \frac{t}{t+s} \sum_{n=0}^{\infty} a_n \overline{b}_n - \frac{s}{t+s} \sum_{n=-\infty}^{-1} a_n \overline{b}_n.$$

Therefore we get a one-parametric family of invariant inner products for $T_{1|0}$. However, all of them are linear combinations of two basis inner products mentioned above ($t = 0$ and $s = 0$).

3 Stein–Sahi representations

Here we extend constructions of the previous section to the groups $G := \mathrm{U}(n,n)$. The analogy of the circle S^1 is the space $\mathrm{U}(n)$ of unitary matrices.

A. Construction of representations

3.1. Distributions $\ell_{\sigma|\tau}$. Let z be an $n \times n$ matrix with norm < 1. For $\sigma \in \mathbb{C}$, we define the function $\det(1-z)^\sigma$ by

$$\det(1-z)^\sigma := \det\Big[1 - \sigma z + \frac{\sigma(\sigma-1)}{2!} z^2 - \frac{\sigma(\sigma-1)(\sigma-2)}{3!} z^3 + \dots\Big].$$

Extend this function to matrices z satisfying $\|z\| \leqslant 1$, $\det(1-z) \neq 0$ by

$$\det(1-z)^\sigma := \lim_{u\to z,\ \|u\|<1} \det(1-u)^\sigma.$$

The expression $\det(1-z)^\sigma$ is continuous in the domain $\|z\| \leqslant 1$ except the surface $\det(1-z) = 0$.

Denote by $\det(1-z)^{\{\sigma|\tau\}}$ the function

$$\det(1-z)^{\{\sigma|\tau\}} := \det(1-z)^\sigma \det(1-\overline{z})^\tau.$$

We define the function $\ell_{\sigma|\tau}(g)$ on the unitary group $\mathrm{U}(n)$ by

$$\ell_{\sigma|\tau}(z) := 2^{-(\sigma+\tau)n} \det(1-z)^{\{\sigma|\tau\}}. \tag{3.1}$$

Obviously,

$$\ell_{\sigma|\tau}(h^{-1}zh) = \ell_{\sigma|\tau}(z) \qquad \text{for } z,\, h \in \mathrm{U}(n). \tag{3.2}$$

Lemma 3.1 *Let* $e^{i\psi_1}$, $\dots$, $e^{i\psi_n}$, *where* $0 \leqslant \psi_k < 2\pi$, *be the eigenvalues of* $z \in \mathrm{U}(n)$. *Then*

$$\ell_{\sigma|\tau}(z) = \exp\Bigl\{\frac{i}{2}(\sigma-\tau)\sum_k(\psi_k-\pi)\Bigr\}\prod_{k=1}^{n}\sin^{\sigma+\tau}\frac{\psi_k}{2}. \tag{3.3}$$

PROOF. It suffices to verify the statement for diagonal matrices; equivalently we must check the identity

$$(1-e^{i\psi})^{\{\sigma|\tau\}} = \exp\Bigl\{\frac{i}{2}(\sigma-\tau)(\psi-\pi)\Bigr\}\sin^{\sigma+\tau}\frac{\psi}{2}.$$

We have

$$\frac{1}{2}(1-e^{i\psi}) = \exp\Bigl\{\frac{i}{2}(\psi-\pi)\Bigr\}\sin\frac{\psi}{2}.$$

Further, both the sides of the equality

$$2^{-\sigma}(1-e^{i\psi})^{\sigma} = \exp\Bigl\{\frac{i}{2}\sigma(\psi-\pi)\Bigr\}\sin^{\sigma}\frac{\psi}{2},$$

are real-analytic on $(0,2\pi)$ and the substitution $\psi=\pi$ gives 1 in both the sides. □

3.2. Positivity. Let $\mathrm{Re}(\sigma+\tau) < 1$. Consider the sesquilinear form on $C^\infty(\mathrm{U}(n))$ given by

$$\langle f_1, f_2\rangle_{\sigma|\tau} = \iint_{\mathrm{U}(n)\times\mathrm{U}(n)} \ell_{\sigma|\tau}(zu^{-1})f_1(z)\overline{f_2(u)}\,d\mu(z)\,d\mu(u). \tag{3.4}$$

For σ, $\tau\in\mathbb{R}$ this form is Hermitian, i.e.,

$$\langle f_2, f_1\rangle_{\sigma|\tau} = \overline{\langle f_1, f_2\rangle}_{\sigma|\tau}$$

Observation 3.2 *For fixed* f_1, $f_2 \in C^\infty(\mathrm{U}(n))$, *this expression admits a meromorphic continuation in* σ, τ *to the whole* $\mathbb{C}^2$.

This follows from general facts about distributions; however, this fact is a corollary of the expansion of the distributions $\ell_{\sigma|\tau}$ in characters, see Theorem 3.11. This expansion implies also the following theorem:

Theorem 3.3 *For* $\sigma,\tau\in\mathbb{R}\setminus\mathbb{Z}$, *the inner product (3.4) is positive definite (up to a sign) iff integer parts of* $-\sigma-n$ *and* τ *are equal.*

The domain of positivity is the union of the dotted squares on Figure 5.

For σ, τ satisfying this theorem, denote by $\mathcal{H}_{\sigma|\tau}$ the completion of $C^\infty(\mathrm{U}(n))$ with respect to our inner product.

3.3. The group $\mathrm{U}(n,n)$**.** Consider the linear space $\mathbb{C}^n\oplus\mathbb{C}^n$ equipped with the indefinite Hermitian form

$$\{v\oplus w, v'\oplus w'\} = \langle v, v'\rangle_{\mathbb{C}^n\oplus 0} - \langle w, w'\rangle_{0\oplus\mathbb{C}^n}, \tag{3.5}$$



1. The dotted squares correspond to unitary representations $\rho_{\sigma|\tau}$.
2. Vertical and horizontal rays in the south-west of Figure correspond to nondegenerate highest weight and lowest weight representations. Fat points correspond to degenerated highest and lowest weight representations, and also to the unipotent representations. The point $(\sigma, \tau) = (0, 0)$ corresponds to the trivial one-dimensional representation.
3. In points of the thick segments, we have some exotic unitary sub-quotients.
4. The shift $(\sigma, \tau) \mapsto (\sigma + 1, \tau - 1)$ send a representation $\rho_{\sigma|\tau}$ of $\mathrm{SU}(n, n)^{\sim}$ to an equivalent representation.
5. The permutation of the axes $(\tau, \sigma) \mapsto (\sigma, \tau)$ gives a complex conjugate representation.
6. The symmetry with respect to the point $(-n/2, -n/2)$ (black circle) gives a dual representation (for odd n this point is a center of a dotted square; for even n this point is a common vertex of two dotted squares).
7. For $\sigma + \tau = n$ (the diagonal line) our Hermitian form is the standard L^2-product.
8. Linear (non-projective) representations of $\mathrm{U}(n, n)$ correspond to the family of parallel lines $\sigma - \tau \in \mathbb{Z}$.

Figure 5: Unitarizability conditions for $\mathrm{U}(n, n)$. The case $n = 5$.

where $\langle\cdot,\cdot\rangle$ is the standard inner product in $\mathbb{C}^n$. Denote by $\mathrm{U}(n,n)$ the group of linear operators in $\mathbb{C}^n\oplus\mathbb{C}^n$ preserving the form $\{\cdot,\cdot\}$. We write elements of this group as block $(n+n)\times(n+n)$ matrices $g:=\begin{pmatrix}a&b\\c&d\end{pmatrix}$. By definition, such matrices satisfy the condition

$$g\begin{pmatrix}1&0\\0&-1\end{pmatrix}g^*=\begin{pmatrix}1&0\\0&-1\end{pmatrix}. \tag{3.6}$$

Lemma 3.4 *The following formula*

$$z\mapsto z^{[g]}:=(a+zc)^{-1}(b+zd),\qquad z\in\mathrm{U}(n),\ g=\begin{pmatrix}a&b\\c&d\end{pmatrix}\in\mathrm{U}(n,n) \tag{3.7}$$

determines an action of the group $\mathrm{U}(n,n)$ *on the* space $\mathrm{U}(n)$.

Proof is given in Subsection 3.20.

3.4. Representations $\rho_{\sigma|\tau}$ of $\mathrm{U}(n,n)$. Denote by $\mathrm{U}(n,n)^\sim$ the universal covering of the group $\mathrm{U}(n,n)$, see for details Subsection 3.18. Fix σ, $\tau\in\mathbb{C}$. We define an action of $\mathrm{U}(n,n)^\sim$ in the space $C^\infty(\mathrm{U}(n))$ by the linear operators

$$\rho_{\sigma|\tau}(g)f(z)=f(z^{[g]})\det{}^{\{-n-\tau|-n-\sigma\}}(a+zc). \tag{3.8}$$

We must explain the meaning of the complex power in this formula. First,

$$a+zc=(1+zca^{-1})a$$

The defining equation (3.6) implies $\|ca^{-1}\|<1$. Hence, for all matrices z satisfying $\|z\|\leqslant 1$, complex powers of $1+zca^{-1}$ are well defined. Next,

$$\det(a)^{-n-\tau|-n-\sigma}:=\exp\Bigl\{-(n+\tau)\ln\det a-(n+\sigma)\overline{\ln\det a}\Bigr\}$$

It is a well-defined function on $\mathrm{U}(n,n)^\sim$. We set

$$\det(a+zc)^{-n-\tau|-n-\sigma}:=\det\Bigl[(1+zca^{-1})^{-n-\tau|-n-\sigma}\Bigr]\det(a)^{-n-\tau|-n-\sigma}$$

3.5. The Stein–Sahi representations.

Proposition 3.5 *The operators $\rho_{\sigma|\tau}(g)$ preserve the form $\langle\cdot,\cdot\rangle_{\sigma|\tau}$.*

Proof is given in Subsection 3.23.

Corollary 3.6 *For σ, τ satisfying the positivity conditions of Theorem 3.3, the representation $\rho_{\sigma|\tau}$ is unitary in the Hilbert space $\mathcal{H}_{\sigma|\tau}$.*

3.6. The degenerate principal series.

Proposition 3.7 *Let* $\mathrm{Re}(\rho+\sigma)=-n$, $\mathrm{Im}\,\sigma=\mathrm{Im}\,\tau$. *Then the representation $\rho_{\sigma|\tau}$ is unitary in $L^2(\mathrm{U}(n))$.*

Figure 6: A '*Maya diagram*' for signatures. We draw the integer 'line' and fill the boxes $m_1, \dots, m_n$ with black.

Proposition 3.8

$$\frac{\langle f_1, f_2\rangle_{\sigma|\tau}}{\prod_{j=1}^{n}\Gamma(\sigma+\tau+j)}\Bigg|_{\sigma=-n-\tau} = \mathrm{const}\cdot\int_{\mathrm{U}(n)} f_1(u)\overline{f_2(u)}\,d\mu(u)$$

.

3.7. Shifts of parameters.

Proposition 3.9 *For integer k,*

$$\rho_{\sigma+k|\tau-k} \simeq (\det g)^k\cdot\rho_{\sigma|\tau}$$

The intertwining operator is the multiplication by the determinant

$$F(z)\mapsto F(z)\det(z)^k.$$

This operator also defines an isometry of the corresponding Hermitian forms.

B. Expansions of distributions $\ell_{\sigma|\tau}$ in characters. Positivity

3.8. Characters of $\mathrm{U}(n)$**.** See Weyl's book [49]. The set of finite dimensional representations of $\mathrm{U}(n)$ is parameterized by collections of integers (*signatures*)

$$\mathbf{m}:\quad m_1 > m_2 > \dots > m_n.$$

The character $\chi_{\mathbf{m}}$ of the representation[6] $\pi_{\mathbf{m}}$ (a *Schur function*) corresponding to a signature $\mathbf{m}$ is given by

$$\chi_{\mathbf{m}}(z) = \frac{\det_{k,j=1,2,\dots,n}\bigl\{e^{im_j\psi_k}\bigr\}}{\det_{k,j=1,2,\dots,n}\bigl\{e^{i(j-1)\psi_k}\bigr\}}, \tag{3.9}$$

where $e^{i\psi_k}$ are the eigenvalues of z. Recall that the denominator admits the decomposition

$$\det_{k,j}\bigl\{e^{i(j-1)\psi_k}\bigr\} = \prod\nolimits_{l<k}(e^{i\psi_l}-e^{i\psi_k}). \tag{3.10}$$

The dimension of π_m is

$$\dim\pi_{\mathbf{m}} = \chi_{\mathbf{m}}(1) = \frac{\prod_{0\leqslant\alpha<\beta\leqslant n}(m_\alpha-m_\beta)}{\prod_{j=1}^{n} j!}. \tag{3.11}$$

---

[6] Explicit constructions of representations of $\mathrm{U}(n)$ are not used below.

3.9. Central functions. A function $F(z)$ on $\mathrm{U}(n)$ is called *central* if

$$F(h^{-1}zh) = F(z) \qquad \text{for all } z,\, h \in \mathrm{U}(n).$$

In particular characters and $\ell_{\sigma|\tau}$ are central functions.

For central functions F on $\mathrm{U}(n)$, the following *Weyl integration formula* holds

$$\int\limits_{\mathrm{U}(n)} F(z)\, d\mu(z) = \frac{1}{(2\pi)^n n!} \int\limits_{0<\psi_1<2\pi} \dots \int\limits_{0<\psi_n<2\pi} F\bigl(\mathrm{diag}(e^{i\psi_1}, \dots, e^{i\psi_n})\bigr) \times \\ \times \Bigl|\prod_{m<k} (e^{i\psi_m} - e^{i\psi_k})\Bigr|^2 \prod_{k=1}^{n} d\varphi_k, \quad (3.12)$$

where $\mathrm{diag}(\cdot)$ is a diagonal matrix with given entries.

Any central function $F \in L^2(\mathrm{U}(n))$ admits an expansion in characters,

$$F(z) = \sum\nolimits_{\mathbf{m}} c_{\mathbf{m}} \chi_m(z),$$

where the summation is given over all signatures $\mathbf{m}$ and the coefficients $c_{\mathbf{m}}$ are L^2-inner products

$$c_{\mathbf{m}} = \int_{\mathrm{U}(n)} F(z) \overline{\chi_{\mathbf{m}}(z)}\, d\mu(z).$$

Note that $\overline{\chi}_{\mathbf{m}} = \chi_{\mathbf{m}^*}$, where

$$\mathbf{m}^* := (n-1-m_n, \dots, n-1-m_2, n-1-m_1)$$

Applying formula (3.12), explicit expression (3.9) for characters, and formula (3.10) for the denominator, we obtain

$$c_{\mathbf{m}} = \frac{1}{(2\pi)^n n!} \int\limits_{0<\psi_1<2\pi} \dots \int\limits_{0<\psi_n<2\pi} F\Bigl(\mathrm{diag}\bigl\{e^{i\psi_1}, \dots, e^{i\psi_n}\bigr\}\Bigr) \times \\ \times \det_{k,j=1,2,\dots,n} \bigl\{e^{i(j-1)\psi_k}\bigr\} \det_{k,j=1,2,\dots,n} \bigl\{e^{-im_j\psi_k}\bigr\} \prod_{k=1}^{n} d\varphi_k. \quad (3.13)$$

Let $F(z)$ be multiplicative with respect to eigenvalues,

$$F(z) = \prod_k f\bigl(e^{i\varphi_k}\bigr)$$

(for, instance $F = \ell_{\sigma|\tau}$, see (3.3)). Then we can apply the following simple lemma (see e.g. [28]).

Lemma 3.10 *Let X be a set,*

$$\int_{X^n} \prod_{k=1}^{n} f(x_k) \det_{k,l=1,\dots n}\{u_l(x_k)\} \det_{k,l=1,\dots n}\{v_l(x_k)\} \prod_{j=1}^{n} dx_j = \\ = n! \det_{l,m=1,\dots,n}\Bigl\{\int_X f(x)u_l(x)v_m(x)\,dx\Bigr\} \qquad (3.14)$$

.

3.10. Lobachevsky beta-integrals. We wish to apply Lemma 3.10 to functions $\ell_{\sigma|\tau}$. For this purpose, we need for the following integral, see [15], 3.631,1, 3.631,8,

$$\int_0^{\pi} \sin^{\mu-1}(\varphi)\, e^{ib\varphi}\, d\varphi = \frac{2^{1-\mu}\pi\Gamma(\mu)e^{ib\pi/2}}{\Gamma\bigl((\mu+b+1)/2\bigr)\Gamma\bigl((\mu-b+1)/2\bigr)}. \qquad (3.15)$$

It is equivalent to the identity (2.12).

In a certain sense, the integral (3.21) is a multivariate analog of the Lobachevsky integral. On the other hand, (3.21) is a special case of the modified Kadell integral [29].

3.11. Expansion of the function $\ell_{\sigma|\tau}$ in characters.

Theorem 3.11 *Let* $\mathrm{Re}(\sigma+\tau)<1$. *Then*

$$\ell_{\sigma|\tau}(g) = \\ = \frac{(-1)^{n(n-1)/2}\sin^n(\pi\sigma)2^{-(\sigma+\tau)n}}{\pi^n}\prod_{j=1}^{n}\Gamma(\sigma+\tau+j)\times \\ \times\sum_{\mathbf{m}}\Bigl\{\prod_{1\leqslant\alpha<\beta\leqslant n}(m_\alpha-m_\beta)\prod_{j=1}^{n}\frac{\Gamma(-\sigma+m_j-n+1)}{\Gamma(\tau+m_j+1)}\chi_{\mathbf{m}}(g)\Bigr\} = \qquad (3.16)$$

$$= (-1)^{n(n-1)/2}2^{-(\sigma+\tau)n}\prod_{j=1}^{n}\Gamma(\sigma+\tau+j)\times \\ \times\sum_{\mathbf{m}}\Biggl\{\frac{(-1)^{\sum m_j}\prod_{1\leqslant\alpha<\beta\leqslant n}(m_\alpha-m_\beta)}{\prod_{j=1}^{n}\Gamma(\sigma-m_j+n)\Gamma(\tau+m_j+1)}\chi_{\mathbf{m}}(g)\Biggr\}. \qquad (3.17)$$

Proof is contained in Subsection 3.13. For the calculation we need for Lemma 3.13 proved in the next subsection.

3.12. A determinant identity. Recall that the *Cauchy determinant* (see e.g. [22]) is given by

$$\det_{kl}\Bigl\{\frac{1}{x_k+y_l}\Bigr\} = \frac{\prod_{1\leqslant k<l\leqslant n}(x_k-x_l)\cdot\prod_{1\leqslant k<l\leqslant n}(y_k-y_l)}{\prod_{1\leqslant k,l\leqslant n}(x_k+y_l)}. \qquad (3.18)$$

The following version of the Cauchy determinant is also well known.

Lemma 3.12

$$\det\begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ \frac{1}{x_1+b_1} & \frac{1}{x_2+b_1} & \frac{1}{x_3+b_1} & \dots & \frac{1}{x_n+b_1} \\ \frac{1}{x_1+b_2} & \frac{1}{x_2+b_2} & \frac{1}{x_3+b_2} & \dots & \frac{1}{x_n+b_2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{1}{x_1+b_{n-1}} & \frac{1}{x_2+b_{n-1}} & \frac{1}{x_3+b_{n-1}} & \dots & \frac{1}{x_n+b_{n-1}} \end{pmatrix} =$$

$$= \frac{\prod_{1\leqslant k<l\leqslant n}(x_k-x_l)\prod_{1\leqslant\alpha<\beta\leqslant n-1}(b_\alpha-b_\beta)}{\prod\limits_{\substack{1\leqslant k\leqslant n\\ 1\leqslant\alpha\leqslant n-1}}(x_k+b_\alpha)}. \quad (3.19)$$

Proof. Let Δ be the Cauchy determinant (3.18). Then

$$y_1\Delta = \begin{pmatrix} \frac{y_1}{x_1+y_1} & \frac{y_1}{x_2+y_1} & \dots & \frac{y_1}{x_1+y_1} \\ \frac{1}{x_1+y_2} & \frac{1}{x_2+y_2} & \dots & \frac{1}{x_n+y_2} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{x_1+y_n} & \frac{1}{x_2+y_n} & \dots & \frac{1}{x_n+y_n} \end{pmatrix}.$$

We take $\lim\limits_{y_1\to\infty} y_1\Delta$ and substitute $y_{\alpha+1}=b_\alpha$. $\square$

The following determinant is a rephrasing of [22], Lemma 3.

Lemma 3.13

$$\det\begin{pmatrix} 1 & 1 & 1 & \dots & 1 \\ \frac{x_1+b_1}{x_1+a_1} & \frac{x_2+a_1}{x_2+b_1} & \frac{x_3+a_1}{x_3+b_1} & \dots & \frac{x_n+a_1}{x_n+b_1} \\ \frac{(x_1+a_1)(x_1+a_2)}{(x_1+b_1)(x_1+b_2)} & \frac{(x_2+a_1)(x_2+a_2)}{(x_2+b_1)(x_2+b_2)} & \frac{(x_3+a_1)(x_3+a_2)}{(x_3+b_1)(x_3+b_2)} & \dots & \frac{(x_n+a_1)(x_n+a_2)}{(x_n+b_1)(x_n+b_2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ \frac{\prod\limits_{1\leqslant m\leqslant n-1}(x_1+a_m)}{\prod\limits_{1\leqslant m\leqslant n-1}(x_1+b_m)} & \frac{\prod\limits_{1\leqslant m\leqslant n-1}(x_2+a_m)}{\prod\limits_{1\leqslant m\leqslant n-1}(x_2+b_m)} & \frac{\prod\limits_{1\leqslant m\leqslant n-1}(x_3+a_m)}{\prod\limits_{1\leqslant m\leqslant n-1}(x_3+b_m)} & \dots & \frac{\prod\limits_{m:\,1\leqslant m\leqslant n-1}(x_n+a_m)}{\prod\limits_{m:\,1\leqslant m\leqslant n-1}(x_n+b_m)} \end{pmatrix} =$$

$$= \frac{\prod\limits_{1\leqslant k<l\leqslant n}(x_k-x_l)\prod\limits_{1\leqslant\alpha\leqslant\beta\leqslant n-1}(a_\alpha-b_\beta)}{\prod\limits_{1\leqslant k\leqslant n,1\leqslant\beta\leqslant n-1}(x_k+b_\beta)}. \quad (3.20)$$

Proof. Decomposing a matrix element into a sum of partial fractions, we obtain

$$\frac{(x_k+a_1)\dots(x_k+a_\alpha)}{(x_k+b_1)\dots(x_k+b_\alpha)} \quad = \quad 1 \ + \sum_{1\leqslant\beta\leqslant\alpha}\frac{\prod_{j\leqslant\alpha}(a_j-b_\beta)}{\prod_{j\leqslant\alpha,j\neq\beta}(b_j-b_\beta)} \ \cdot \ \frac{1}{x_k+b_\beta}$$

Therefore the $(\alpha+1)$-th row is a linear combination of the following rows:

$$\begin{array}{cccccc} \Bigl(& 1 & 1 & \dots & 1 & \Bigr), \\ \Bigl(& \frac{1}{x_1+b_1} & \frac{1}{x_2+b_1} & \dots & \frac{1}{x_n+b_1} & \Bigr), \\ \dots & \dots & \dots & \dots & \dots & \dots \\ \Bigl(& \frac{1}{x_1+b_\alpha} & \frac{1}{x_2+b_\alpha} & \dots & \frac{1}{x_n+b_\alpha} & \Bigr). \end{array}$$

Thus our determinant equals

$$\prod_{\alpha=1}^{l-1} \frac{\prod_{j=1}^{\alpha}(a_j-b_\alpha)}{\prod_{j=1}^{\alpha-1}(b_j-b_\alpha)} \cdot \det \begin{pmatrix} 1 & 1 & \dots & 1 \\ \frac{1}{x_1+b_1} & \frac{1}{x_2+b_1} & \dots & \frac{1}{x_n+b_1} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{1}{x_1+b_\alpha} & \frac{1}{x_2+b_\alpha} & \dots & \frac{1}{x_n+b_\alpha} \end{pmatrix}.$$

and we refer to Lemma 3.12. □

3.13. Proof of Theorem 3.11. We must evaluate the inner product

$$\int_{\mathrm{U}(n)} \ell_{\sigma|\tau}(g)\,\overline{\chi_{\mathbf{m}}(g)}\,d\mu(g).$$

Applying (3.13), we get

$$\frac{1}{(2\pi)^n\,n!} \int\limits_{0<\psi_k<2\pi} \prod_{j=1}^{n} \Bigl[\sin^{\sigma+\tau}\bigl(\psi_j/2\bigr)\cdot \exp\Bigl\{\frac{i}{2}(\sigma-\tau)(\psi_j-\pi)\Bigr\}\Bigr]\times$$

$$\times \det_{1\leqslant k,l\leqslant n}\{e^{-im_k\psi_l}\}\cdot \det_{1\leqslant k,l\leqslant n}\{e^{i(k-1)\psi_l}\}\prod_{l=1}^{n} d\psi_l. \qquad (3.21)$$

By Lemma 3.10, we reduce this integral to

$$\frac{1}{(2\pi)^n}\det_{1\leqslant k,j\leqslant n} I(k,j),$$

where

$$I(k,j)=e^{-i(\sigma-\tau)\pi/2}\int_0^{2\pi} \sin^{\sigma-\tau}(\psi/2)\cdot \exp\bigl\{i(\,(\sigma+\tau)/2+k-1-m_j)\bigr\}\,d\psi.$$

We apply the Lobachevsky integral (3.15) and get

$$I(k,j)=\frac{2^{1-\sigma-\tau}\pi\Gamma(\sigma+\tau+1)\,(-1)^{k-1-m_j}}{\Gamma(\sigma+k-m_j)\Gamma(\tau-k+m_j+2)}$$

Applying standard formulas for Γ-function, we come to

$$I(k,j)=2^{1-\sigma-\tau}\Gamma(\sigma+\tau+1)\,\sin(-\sigma\pi)\cdot\frac{\Gamma(-\sigma+m_j-k+1)}{\Gamma(\tau+m_j-k+2)}=$$

$$=2^{1-\sigma-\tau}\Gamma(\sigma+\tau+1)\,\sin(-\sigma\pi)\cdot\frac{\Gamma(-\sigma+m_j-n+1)}{\Gamma(\tau+m_j-n+2)}\cdot\boxed{\frac{(-\sigma+m_j-n+1)_{n-k}}{(\tau+m_j-n+2)_{n-k}}}$$

The factors outside the box do not depend on on k. Thus, we must evaluate the determinant

$$\det_{1\leqslant k,j\leqslant n} \frac{(-\sigma+m_j-n+1)_{n-k}}{(\tau+m_j-n+2)_{n-k}}.$$

Up to a permutation of rows, it is a determinant of the form described in Lemma 3.13 with

$$x_j = m_j, \qquad a_j = -\sigma - n + j, \qquad b = \tau - n + j + 1.$$

After a rearrangement of the factors, we obtain the required result. $\square$

3.14. Characters of compact groups. Preliminaries. First, recall some standard facts on characters of compact groups, for details, see e.g. [21], 9.2, 11.1.

Let K be a compact Lie group equipped with the Haar measure μ, let $\mu(K) = 1$. Let π_1, π_2, …be the complete collection of pairwise distinct irreducible representations of K. Let χ_1, χ_2, …be their characters. Recall the orthogonality relations,

$$\langle \chi_k, \chi_l\rangle_{L^2(K)} = \int_K \chi_k(h)\overline{\chi_l(h)}\, d\mu(h) = \delta_{k,l} \tag{3.22}$$

and

$$\chi_k * \chi_l = \begin{cases} \frac{1}{\dim \pi_k}\chi_k & \text{if } k = l,\\ 0 & \text{if } k \neq l.\end{cases} \tag{3.23}$$

where $*$ denotes the convolution on the group,

$$u * v(g) = \int_K u(gh^{-1})\, v(h)\, d\mu(h).$$

Consider the action of the group $K \times K$ in $L^2(K)$ by the left and right shifts

$$(k_1, k_2):\ f(g) \mapsto f(k_1^{-1} g k_2).$$

The representation of $K \times K$ in $L^2(K)$ is a multiplicity free direct sum of irreducible representations having the form $\pi_k \otimes \pi_k^*$, where π_k^* denotes the dual representation,

$$L^2(K) \simeq \bigoplus_k \pi_k \otimes \pi_k^*. \tag{3.24}$$

Denote by $V_k \subset L^2(K)$ the space of representation $\pi_k \otimes \pi_k^*$. Each distribution f on K is a sum of 'elementary harmonics',

$$f = \sum\nolimits_k f^k, \qquad f_k \in V_k..$$

The projector to a subspace V_k is the convolution with the corresponding character,

$$f^k = \frac{1}{\dim \pi_k} f * \chi_k \tag{3.25}$$

(in particular, f^k is smooth).

Observation 3.14 *Let f be a function on* $\mathrm{U}(n)$, $f = \sum_{\mathbf{m}} a_m f^{\mathbf{m}}$, *where* $f^{\mathbf{m}} \in V_{\mathbf{m}}$.

a) $f \in C^\infty(\mathrm{U}(n))$ *iff*

$$\|f^{\mathbf{m}}\|_{L^2} = o\left(\sum m_j^2\right)^{-L} \qquad \text{for all } L.$$

b) f *is a distribution on* $\mathrm{U}(n)$ *iff there exists* L *such that*

$$\|f^{\mathbf{m}}\|_{L^2} = o\left(\sum m_j^2\right)^{L}.$$

Proof: Note that $f \in L^2(\mathrm{U}(n))$ iff $\sum \|f^{\mathbf{m}}\|_{L^2}^2 < \infty$. Denote by Δ be the second order invariant Laplace operator on $\mathrm{U}(n)$. Then $\Delta f^{\mathbf{m}} = q(\mathbf{m}) f^{\mathbf{m}}$, where $q(\mathbf{m}) = \sum m_j^2 + \dots$ is an explicit quadratic expression in $\mathbf{m}$. For $f \in C^\infty$ we have $\Delta^p f \in C^\infty$; this implies the first statement. Since $q(\mathbf{m})$ has a finite number of zeros (one), the second statement follows from a) and the duality. $\square$

3.15. Hermitian forms defined by kernels. Let Ξ be a central distribution on K satisfying $\Xi(g^{-1}) = \overline{\Xi(g)}$. Consider the following Hermitian form on $C^\infty(K)$

$$\langle f_1, f_2 \rangle = \iint_{K\times K} \Xi(gh^{-1})\, f_1(h)\overline{f_2(g)}\, d\mu(h)\, d\mu(g). \tag{3.26}$$

Consider the expansion of Ξ in characters

$$\Xi = \sum_k c_k \chi_k.$$

Lemma 3.15

$$\langle f_1, f_2 \rangle = \sum_k \frac{c_k}{\dim \pi_k} \int_{\mathrm{U}(n)} f_1^k(h)\overline{f_2^k(h)}\, d\mu(h). \tag{3.27}$$

Proof. The Hermitian form (3.26) is $K \times K$-invariant. Therefore the subspaces $V_k \simeq \pi_k \otimes \pi_k^*$ must be pairwise orthogonal. Since $\pi_k \otimes \pi_k^*$ is an irreducible representation of $K \times K$, it admits a unique up to a factor $K \times K$-invariant Hermitian form. Therefore it is sufficient to find these factors.

Set $f_1 = f_2 = \chi_k$. We evaluate

$$\iint_{K\times K} \left(\sum_k c_k \chi_k(gh^{-1})\right) \chi_k(h)\overline{\chi_k(g)}\, d\mu(g)\, d\mu(h) = \frac{c_k}{\dim \pi_k}$$

using (3.22) and (3.23). $\square$

3.16. Positivity. Let $\mathrm{Re}(\sigma + \tau) < 1$. Consider the sesquilinear form on $C^\infty(\mathrm{U}(n))$ given by

$$\langle f_1, f_2 \rangle_{\sigma|\tau} = \iint_{\mathrm{U}(n)\times \mathrm{U}(n)} \ell_{\sigma|\tau}(zu^{-1}) f_1(z)\overline{f_2(u)}\, d\mu(z)\, d\mu(u), \tag{3.28}$$

where the distribution $\ell_{\sigma|\tau}$ is the same as above.

Observation 3.16 *For fixed f_1, $f_2 \in C^\infty(\mathrm{U}(n))$, the expression $\langle f_1, f_2\rangle_{\sigma|\tau}$ admits a meromorphic continuation in σ, τ to the whole $\mathbb{C}^2$.*

Proof. Expanding f_1, f_2 in elementary harmonics

$$f_1(z) = \sum_{\mathbf{m}} f_2^{\mathbf{m}}(z), \qquad f_2(z) = \sum_{\mathbf{m}} f_2^{\mathbf{m}}(z),$$

we get (see Lemma 3.15)

$$\langle f_1, f_2\rangle_{\sigma|\tau} = \sum_{\mathbf{m}} \frac{c_{\mathbf{m}}}{\dim \pi_{\mathbf{m}}} \int_{\mathrm{U}(n)} f_1^{\mathbf{m}}(z)\overline{f_2^{\mathbf{m}}(z)}\, d\mu(z),$$

where the meromorphic expressions for $c_{\mathbf{m}}$ were obtained in Theorem 3.11. The coefficients $c_{\mathbf{m}}$ have polynomial growth in $\mathbf{m}$. On the other hand, $\|f_j^{\mathbf{m}}\|$ rapidly decrease, see Observation 3.14. Therefore, the series converges. $\square$

Proof of positivity. Corollary 3.6 We look at the expression (3.16). It suffices to examine the factor

$$\frac{\Gamma(-\sigma - n + m_j + 1)}{\Gamma(\tau + m_j + 1)}, \tag{3.29}$$

because signs of all the remaining factors are independent on m_j. Let $n \in \mathbb{Z}$ and $\alpha \in (0, 1)$. Then

$$\operatorname{sign} \Gamma(n + \alpha) = \begin{cases} +1, & \text{if } n \geqslant> 0, \\ (-1)^n, & \text{if } n < 0 \end{cases}$$

Therefore (3.29) is positive whenever integer parts of τ and $-\sigma - n$ equal. $\square$

3.17. The L^2-limit. Proof of Proposition 3.8. Thus, let $\sigma + \tau = -n$. Then

$$\Bigl(\prod_{j=1}^n \Gamma(\sigma + \tau + j)\Bigr)^{-1} \ell_{\sigma|\tau} = \text{const} \cdot \sum (\dim \pi_{\mathbf{m}}) \chi_{\mathbf{m}}$$

Indeed, in this case Γ-factors in (3.16) cancel, and we use (3.11).

Keeping in mind (3.27), we get Proposition 3.8.

C. Other proofs

Here we prove that the operators $\rho_{\sigma|\tau}$ preserve the inner product determined by the distribution $\ell_{\sigma|\tau}$.

3.18. The universal covering of the group U(n, n). The fundamental group of U(n, n) is[7]

$$\pi_1\bigl(\mathrm{U}(n, n)\bigr) \simeq \mathbb{Z} \oplus \mathbb{Z}.$$

---

[7] By a general theorem, a real reductive Lie group G admits a deformation retraction to its maximal compact subgroup K. In our case, $K = \mathrm{U}(n) \times \mathrm{U}(n)$ and $\pi_1(\mathrm{U}(n)) = \mathbb{Z}$.

The universal covering $\mathrm{U}(n,n)^\sim$ of $\mathrm{U}(n,n)$ can be identified with the the set $\mathfrak{U}$ of triples

$$\left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix},\, s,\, t \right\} \in \mathrm{U}(n,n) \times \mathbb{C} \times \mathbb{C}$$

satisfying the conditions

$$\det(a) = e^s, \qquad \det(d) = e^t.$$

The multiplication of triples is given by the formula

$$(g_1, s_1, t_1) \circ (g_2, s_2, t_2) = \big(g_1 g_2, s_1 + s_2 + c^+(g_1, g_2), t_1 + t_2 + c^-(g_1, g_2)\big),$$

where the *Berezin cocycle* $c^\pm$ is given by

$$c^+(g_1, g_2) = \operatorname{tr} \ln(a_1^{-1} a_3 a_2^{-1}), \qquad c^-(g_1, g_2) = \operatorname{tr} \ln(d_1^{-1} d_3 d_2^{-1});$$

here $g_3 = g_1 g_2$, and $g_j = \begin{pmatrix} a_j & b_j \\ c_j & d_j \end{pmatrix}$. It can be shown that $\|a_1^{-1} a_3 a_2^{-1} - 1\| < 1$, therefore the logarithm is well defined. On the other hand,

$$e^{s_3} = e^{s_1 + s_2 + c^+(g_1, g_2)} = \det(a_1)\det(a_2)\det(a_1^{-1} a_3 a_2^{-1}) = \det(a_3)$$

This shows that the $\mathfrak{U}$ is closed with respect to multiplication.

For details, see [31].

In particular, $\det(a)$ is a well-defined single-valued function on $\mathrm{U}(n,n)^\sim$. In our notation, it is given by

$$(g, s, t) \mapsto s.$$

3.19. Another model of $\mathrm{U}(n,n)$. We can realize $\mathrm{U}(n,n)$ as the group of $(n+n) \times (n+n)$-matrices $g = \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix}$ satisfying the condition

$$g \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} g^* = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix} \tag{3.30}$$

3.20. Action of $\mathrm{U}(n,n)$ on the space $\mathrm{U}(n)$. Proof of Lemma 3.4 . We must show that for

$$z \in \mathrm{U}(n) \quad \text{and} \quad g = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{U}(n,n)$$

we have

$$z^{[g]} := (a + zc)^{-1}(b + zd) \in \mathrm{U}(n,n) \tag{3.31}$$

For $z \in \mathrm{U}(n)$, consider its graph $graph(z) \subset \mathbb{C}^n \oplus \mathbb{C}^n$. It is an n-dimensional linear subspace, consisting of all vectors $v \oplus vz$, where a vector-row v ranges in $\mathbb{C}^n$. Since $z \in \mathrm{U}(n)$, the subspace $graph(z)$ is isotropic[8] with respect to the

---

[8] A subspace V in a linear space is *isotropic* with respect to an Hermitian form Q if Q equals 0 on V.

Hermitian form $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Conversely, any n-dimensional isotropic subspace in $\mathbb{C}^n \oplus \mathbb{C}^n$ is a graph of a unitary operator $z \in \mathrm{U}(n)$.

Thus we get a one-to-one correspondence between the group $\mathrm{U}(n)$ and the Grassmannian of n-dimensional isotropic subspaces in $\mathbb{C}^n \oplus \mathbb{C}^n$.

The group $\mathrm{U}(n,n)$ acts on the Grassmannian and therefore $\mathrm{U}(n,n)$ acts on the space $\mathrm{U}(n)$. Then (3.31) is the explicit expression for the latter action. Indeed

$$\begin{pmatrix} v \oplus vz \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} = v(a+zc) \,\oplus\, v(b+zd)$$

We denote $\xi := v(a+zc)$ and get

$$\xi \,\oplus\, \xi(a+zc)^{-1}(b+zd)$$

and this completes the proof of Lemma 3.4. $\square$

Thus, $\mathrm{U}(n)$ is a $\mathrm{U}(n,n)$-homogeneous space. We describe without proof (it is a simple exercise) the stabilizer of a point $z = 1$. It is a maximal parabolic subgroup.

In the model (3.30) it can be realized as the subgroup of matrices having the structure

$$\begin{pmatrix} \alpha & 0 \\ \beta & \alpha^{*-1} \end{pmatrix}$$

It is a semidirect product of $\mathrm{GL}(n,\mathbb{C})$ and the Abelian group $\mathbb{R}^{n^2}$.

In our basic model the stabilizer of $z = 1$ is the semi-direct product of two subgroups

$$\frac{1}{2}\begin{pmatrix} \alpha + \alpha^{*-1} & \alpha - \alpha^{*-1} \\ \alpha - \alpha^{*-1} & \alpha + \alpha^{*-1}, \end{pmatrix} \quad \text{where } g \in \mathrm{GL}(n,\mathbb{C}), \tag{3.32}$$

and

$$\begin{pmatrix} 1+iT & iT \\ -iT & 1-iT \end{pmatrix}, \quad \text{where } T = T^*. \tag{3.33}$$

3.21. The Jacobian.

Lemma 3.17 *For the Haar measure $\mu(z)$ on $\mathrm{U}(n)$, we have*

$$\mu\big(z^{[g]}\big) = |\det{}^{-2n}(a+zc)| \cdot \mu(z). \tag{3.34}$$

PROOF. A verification of this formula is straightforward, we only outline the main steps. First, $J(g,z) := |\det{}^{-2n}(a+zc)|$ satisfies the chain rule (1.3). Next, the formula (3.34) is valid for $g \in \mathrm{U}(n,n)$ having the form $\begin{pmatrix} a & 0 \\ 0 & d \end{pmatrix}$, where u, $v \in \mathrm{U}(n)$. Indeed, the corresponding transformation of $u \mapsto u^{[h]}$ is $u \mapsto a^{-1}ud$, its Jacobian is 1.

Therefore we can set $z = 1$, $z^{[g]} = 1$. Now we must evaluate the determinants of the differentials of maps $z \mapsto z^{[g]}$ at $z = 1$ for g given by (3.32) and (3.33). In the second case the differential is the identity map, in the first case the differential is $dz \mapsto \alpha^*(dz)\alpha$. We represent α as $p\Delta q$, where Δ is diagonal with real eigenvalues and p, q are unitary. Now the statement becomes obvious. $\square$

3.22. The degenerate principal series. Proof of Proposition 3.7. Thus, let $\mathrm{Re}(\sigma+\tau)=-n$, $\mathrm{Im}(\sigma)=\mathrm{Im}(\tau)=s$. Then

$$\det(a+uc)^{-n-\sigma|-n-\tau} = |\det(a+uc)|^{-n-2is} e^{i(\tau-\sigma)\mathrm{Arg}\det(a+uc)},$$

where $\mathrm{Arg}(\cdot)$ is the argument of a complex number. Therefore

$$\langle T_{\sigma|\tau}(g)f_1, T_{\sigma|\tau}(g)f_2\rangle_{L^2(\mathrm{U}(n))} = \\ = \int_{\mathrm{U}(n)} f_1(u^{[g]})\overline{f_2(u^{[g]})}\left|\det(a+uc)^{-n-\sigma|-n-\tau}\right|^2 d\mu(u) = \\ = \int_{\mathrm{U}(n)} f_1(u^{[g]})\overline{f_2(u^{[g]})}|\det(a+uc)|^{-2n}\, d\mu(u)$$

and we change the variable $z=u^{[g]}$ keeping in mind Lemma 3.17. $\square$

3.23. The invariance of the kernel. Proof of Proposition 3.5.

Lemma 3.18 *The distribution $\ell_{\sigma|\tau}$ satisfies the identity*

$$\ell_{\sigma|\tau}(u^{[g]}(v^{[g]})^*) = \ell_{\sigma|\tau}(uv^*)\det(a+uc)^{\{-\tau|-\sigma\}}\det(a+vc)^{\{-\sigma|-\tau\}}. \tag{3.35}$$

Proof. This follows from the identity

$$1-u^{[g]}(v^{[g]})^* = (a+uc)^{-1}(1-uv^*)(a+vc)^{*-1}, \qquad \text{where } g\in\mathrm{U}(n,n),$$

which can be easily verified by a straightforward calculation (see e.g. [31]). $\square$

Proof of Proposition 3.5. First, let $\mathrm{Re}(\sigma+\tau)<1$. Substitute $h_1=u_1^{[g]}$, $h_2=u_2^{[g]}$ to the integral

$$\langle f_1,f_2\rangle_{\sigma|\tau} = \iint_{\mathrm{U}(n)\times\mathrm{U}(n)} \ell_{\sigma|\tau}(h_1h_2^*)\, f_1(h_1)\overline{f_2(h_2)}\, d\mu(h_1)\, d\mu(h_2).$$

By the lemma, we obtain

$$\iint_{\mathrm{U}(n)\times\mathrm{U}(n)} \ell_{\sigma|\tau}(u_1u_2^*)\det(a+u_1c)^{\{-\tau|-\sigma\}}|\det(a+u_2c)|^{-\sigma|-\tau\}}\times \\ \times\ f_1(u_1)\overline{f_2(u_2)}|\det(a+u_1c)|^{-2n}|\det(a+u_2c)|^{-2n}\, d\mu(u_1)\, d\mu(u_2) = \\ = \langle \rho_{\sigma|\tau}(g)f_1, \rho_{\sigma|\tau}(g)f_2\rangle_{\sigma|\tau}.$$

Thus, our operators preserve the form $\langle\cdot,\cdot\rangle_{\sigma|\tau}$.

For general σ, $\tau\in\mathbb{C}$, we consider the analytic continuation. $\square$

3.24. Shift of parameters. Proof of Proposition 3.9. First, we recall *Cartan decomposition.* For $t_1\geqslant\ldots\geqslant t_n$ denote

$$\mathrm{CH}(t) := \begin{pmatrix} \cosh(t_1) & 0 & \ldots \\ 0 & \cosh(t_2) & \ldots \\ \vdots & \vdots & \ddots \end{pmatrix}, \quad \mathrm{SH}(t) := \begin{pmatrix} \sinh(t_1) & 0 & \ldots \\ 0 & \sinh(t_2) & \ldots \\ \vdots & \vdots & \ddots \end{pmatrix}.$$

The following statement is well known



Figure 7: Conditions of positivity of holomorphic representations ξ_σ (the '*Berezin–Wallach set*').

Proposition 3.19 *Each element* $g \in \mathrm{U}(n,n)$ *can be represented in the form*

$$g = \begin{pmatrix} u_1 & 0 \\ 0 & v_1 \end{pmatrix} \begin{pmatrix} \mathrm{CH}(t) & \mathrm{SH}(t) \\ \mathrm{SH}(t) & \mathrm{CH}(t) \end{pmatrix} \begin{pmatrix} u_2 & 0 \\ 0 & v_2 \end{pmatrix} \tag{3.36}$$

for some (uniquely determined) t *and some* u_1, u_2, v_1, $v_2 \in \mathrm{U}(n)$.

Now we must show that the operator $f(z) \mapsto \det(z) f(z)$ intertwines $\rho_{\sigma|\tau}$ and $\rho_{\sigma+1|\tau-1}$ A straightforward calculation reduces this to the identity

$$\frac{\det(a+zc)}{\det \overline{(a+zc)}} = \frac{\det(z^{[g]})}{\det(z)},$$

which becomes obvious after the substitution (3.36).

Also,

$$\ell_{\sigma+1|\tau-1}(z) = -\ell_{\sigma|\tau}(z) \det z$$

and this easily implies the second statement of Proposition 3.9.

4 Hilbert spaces of holomorphic functions

Theorem 3.3 exhaust the cases when the form $\langle \cdot, \cdot \rangle_{\sigma|\tau}$ is positive definite on $C^\infty(\mathrm{U}(n))$. However there are cases of positive semi-definiteness. They are discussed in the next two sections.

Set $\tau = 0$. In this case, our construction produces holomorphic representations[9] of $\mathrm{U}(n,n)$. Holomorphic representations were discovered by Harish-Chandra (holomorphic discrete series, [17]) and Berezin (analytic continuations of holomorphic discrete series, [3]). They are discussed in numerous texts (for partial expositions and further references, see e.g. [10], [31]), our aim is to show a link with our considerations.

4.1. The case $\tau = 0$. Substituting $\tau = 0$, we get the action

$$\rho_{\sigma|0}(g)\, f(z) = f(z^{[g]}) \det(a+zc)^{-n} \overline{\det(a+zc)}^{\,-n-\sigma}.$$

The Hermitian form is

$$\langle f_1, f_2 \rangle_{\sigma|0} = \int_{\mathrm{U}(n)} \int_{\mathrm{U}(n)} \det(1 - z^* u)^\sigma f_1(z)\, \overline{f_2(u)}\, d\mu(z)\, d\mu(u).$$

---

[9] or highest weight representations,

Theorem 4.1 *The form* $\langle f_1, f_2\rangle_{\sigma|0}$ *is positive semi-definite iff* σ *is contained in the set*

$$\sigma = 0,\ -1,\ \dots,\ -(n-1),\ \text{or}\ \sigma < -(n-1).$$

This means that all coefficients $c_{\mathbf{m}}$ in the formula (3.27) are non-negative, but some coefficients vanish. In fact the proof (see below) is the examination of these coefficients.

Under the conditions of the theorem we get a structure of a pre-Hilbert space in $C^\infty(\mathrm{U}(n))$. Denote by $\mathcal{H}_\sigma$ the corresponding Hilbert space.

Next, consider the action of the subgroup $\mathrm{U}(n)\times\mathrm{U}(n)$ in $\mathcal{H}_\sigma$. We must get an orthogonal direct sum

$$\bigoplus_{\mathbf{m}\in\Omega_\sigma} \pi_{\mathbf{m}} \oplus \pi^*_{\mathbf{m}}$$

Some of summands of (3.24) disappear, when we pass to the quotient space; actually the summation is taken over a proper subset Ω_σ of the set of all representations. The next theorem is the description of of the set Ω_σ.

Theorem 4.2 *a) If* $\sigma < -(n-1)$, *then*

$$\Omega_\sigma := \Bigl\{\mathbf{m} :\ m_n \geqslant 0\Bigr\}.$$

b) If $\sigma = -n+\alpha$, *where* $\alpha = 1,\ 2,\ \dots,\ n-1,\ n$, *then*

$$\Omega_\sigma = \Bigl\{\mathbf{m} : m_n = 0,\ m_{n-1} = 1,\ \dots,\ m_{n-\alpha+1} = \alpha - 1\Bigr\}.$$

Proofs.[10] Substitute $\tau = 0$ to (3.17),

$$c_{\mathbf{m}} = (-1)^{n(n-1)/2} 2^{-\sigma n} \prod_{j=1}^{n} \Gamma(\sigma + j)\times$$
$$\times \sum_{\mathbf{m}} \left\{ \frac{(-1)^{\sum m_j} \prod\limits_{1\leqslant\alpha<\beta\leqslant n} (m_\alpha - m_\beta)}{\prod\limits_{j=1}^{n} \Gamma(\sigma - m_j + n)\boxed{\Gamma(m_j+1)}} \chi_{\mathbf{m}}(g) \right\} = \qquad (4.1)$$

$$= \frac{(-1)^{n(n-1)/2} \sin^n(\pi\sigma) 2^{-(\sigma)n}}{\pi^n} \prod_{j=1}^{n} \Gamma(\sigma + j)\times \qquad (4.2)$$

$$\times \sum_{\mathbf{m}} \left\{ \prod_{1\leqslant\alpha<\beta\leqslant n} (m_\alpha - m_\beta) \prod_{j=1}^{n} \frac{\Gamma(-\sigma + m_j - n + 1)}{\boxed{\Gamma(m_j+1)}} \chi_{\mathbf{m}}(g) \right\} \qquad (4.3)$$

---

[10] This is the original Berezin's proof, he started from explicit expansions of reproducing kernels (4.6).



Figure 8: 'Maya diagrams' for signatures of harmonics in holomorphic representations.
a) A general case, $\sigma < n-1$.
b) Degenerate case. Here $\sigma = -(n-1)+5$.

We have $\Gamma(m_j+1)=\infty$ for $m_j<0$. Therefore the corresponding fractions in (4.3) are zero, and the expansion of $\ell_{\sigma|0}$ has the form

$$\ell_{\sigma|0} = \sum_{\mathbf{m}\colon m_n \geqslant 0} c_{\mathbf{m}} \chi_{\mathbf{m}}. \tag{4.4}$$

Let us list possible cases.

Case 1. If $\sigma < -n-1$, then all coefficients $c_{\mathbf{m}}$ are positive, see (4.3); in the line (4.2) poles of the Gamma-functions cancel with zeros of sines.

Case 2. If $\sigma \geqslant -n-1$ is non-integer, then all the coefficients $c_{\mathbf{m}}$ are non-zero, but they have different signs.

Case 3. Let σ be integer, $\sigma \geqslant -n+1$. Consider a small perturbation of σ,

$$\sigma = -n+\alpha+\varepsilon.$$

In this case we get an uncertainty in the expression (4.1):

$$\frac{\prod_{j=1}^n \Gamma(-n+\alpha+\varepsilon+j)}{\prod_{j=1}^n \Gamma(\alpha-m_j+\varepsilon)}, \qquad \varepsilon \to 0.$$

The order of the pole of the numerator is $n-\alpha$. However order of a pole in the denominator ranges between $n-\alpha$ and n according to $\mathbf{m}$. If the last order $> n-\alpha$, then the ratio is zero. The only possibility to get order of a pole $= n-\alpha$ is to set

$$m_n = 0, \quad m_{n-1} = 1, \quad \ldots, \quad m_{n-\alpha+1} = 0. \tag{4.5}$$

Thus the coefficients $c_{\mathbf{m}}$ are nonzero only for signatures satisfying (4.5); they are positive.

We omit a discussion of positive integer σ (the invariant inner product is not positive). $\square$

4.2. Intertwining operators. Denote by B_n the space of complex $n \times n$-matrices with norm < 1.

Consider the integral operator

$$I_\sigma f(z) = \int_{\mathrm{U}(n)} \det(1 - zh^*)^\sigma f(h)\, d\mu(h), \qquad z \in \mathrm{B}_n.$$

It intertwines $\rho_{\sigma|0}$ with the representation $\rho_{-n|-n-\sigma}$, denote the last representation by ξ_σ

$$\xi_\sigma(g) f(z) = f(z^{[g]}) \det(a + zc)^\sigma.$$

The I_σ-image $\mathcal{H}^\circ_\sigma$ of the space $\mathcal{H}_\sigma$ consists of functions holomorphic in B_n. The structure of a Hilbert space in the space of holomorphic functions is determined by the reproducing kernel

$$K_\alpha(z, \overline{u}) = \det(1 - zu^*)^\sigma. \tag{4.6}$$

4.3. Concluding remarks (without proofs).

a) For $\sigma < -(2n-1)$, the inner product in $\mathcal{H}^\circ_\sigma$ can be written as an integral

$$\langle f_1, f_2 \rangle = \mathrm{const} \int_{\mathrm{B}_n} f_1(z)\, \overline{f_2(z)}\, \det(1 - zz^*)^{-\sigma-2n}\, dz\, \overline{dz}.$$

b) For $\sigma < n - 1$ the space $\mathcal{H}^\circ_\sigma$ contains all polynomials.

c) Let $\sigma = 0$, -1, …,$-(n-1)$. Consider the matrix

$$\Delta = \begin{pmatrix} \frac{\partial}{\partial z_{11}} & \cdots & \frac{\partial}{\partial z_{1n}} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial z_{n1}} & \cdots & \frac{\partial}{\partial z_{nn}} \end{pmatrix}.$$

The space $\mathcal{H}^\circ_{-(n-1)}$ consists of functions f satisfying the partial differential equation

$$(\det \Delta) f(z) = 0.$$

The space $\mathcal{H}^\circ_\sigma$, where $\sigma = 0$, -1, …, $-(n-1)$, consists of functions that are annihilated by all $(-\sigma+1) \times (-\sigma+1)$-minors of the matrix Δ. Also, $\mathcal{H}^\circ_\sigma$ contains all polynomial satisfying this system of equations.

In particular, the space $\mathcal{H}^\circ_0$ is one-dimensional.

5 Unipotent representations.

Here we propose models for 'unipotent representations' of Sahi [43] and Dvorsky–Sahi, [8]–[9].



Figure 9: Maya diagram for signatures $\in Z_j$; here j is the number of black boxes at the left of the 'obligatory part'.



Figure 10: The case $n = 2$.
a) $\alpha = 0$. The decomposition of $L^2(\mathrm{U}(2))$ into a direct sum.
b) $\alpha = 1$. White circles correspond to the big subrepresentation W_{tail}. The quotient is a direct sum of two subrepresentations.
c) $\alpha = 2$. The quotient is one-dimensional.
d) $0 < \tau < 1$, $\sigma = -n$. The invariant filtration. The subquotients are unitary.

5.1. Quotients of $\rho_{\sigma|\tau}$ at integer points. Set

$$\tau = 0, \qquad \sigma = -n + \alpha, \qquad \text{where } \alpha = 0,\, 1,\, \dots,\, n-1. \tag{5.1}$$

For $j = 0,\, 1,\, \dots,\, n-\alpha$ denote by Z_j the set of all signatures $\mathbf{m}$ of the form

$$\mathbf{m} = (m_1, \dots, m_{n-\alpha-j}, \alpha-1, \alpha-2, \dots, 0, m_{n-j+1}, \dots m_n)$$

Denote by $V_{\mathbf{m}}$ the $\mathrm{U}(n)\times\mathrm{U}(n)$-subrepresentation in $C^\infty(\mathrm{U}(n))$ corresponding a signature $\mathbf{m}$, see Subsection 3.14.

Theorem 5.1 *The subspace*

$$W_{tail} := \bigoplus_{\mathbf{m}\notin\cup Z_j} V_{\mathbf{m}} \subset C^\infty(\mathrm{U}(n)),$$

is $\mathrm{U}(n,n)$*-invariant.*

b) The quotient $C^\infty(\mathrm{U}(n))/W_{tail}$ *is a sum* $n-\alpha+1$ *subrepresentations*

$$W_j = \oplus_{\mathbf{m}\in Z_j} V_{\mathbf{m}}.$$

The representation of $\mathrm{U}(n,n)$ *in each* W_j *is unitary.*

We formulate the result for $\alpha = 0$ separately. In this case $W_{tail} = 0$.

Theorem 5.2 *The representation* $\rho_{-n|0}$ *is a direct sum of* $n+1$ *unitary representations* W_j*, where* $0 \leqslant j \leqslant n$*. We have* $V_{\mathbf{m}} \subset W_j$ *if the number of negative labels* m_k *is* j*.*

In particular, *we get a canonical decomposition of* $L^2(\mathrm{U}(n))$ *into a direct sum of* $(n+1)$ *subspaces.*

Proof is given in the next subsection.

5.2. The blow-up construction. [11] The distribution $\ell_{\sigma|\tau}$ depends meromorphically in two complex variables σ, τ. Its poles and zeros are located at $\sigma\in\mathbb{Z}$ and in $\tau\in\mathbb{Z}$. For this reason, values of $\ell_{\sigma|\tau}$ at points $(\sigma,\tau)\in\mathbb{Z}^2$ generally are not uniquely defined. Passing to such points from different directions, we get different limits[12].

Thus, set

$$\sigma = -n + \alpha + s\varepsilon, \quad \tau = t\varepsilon \qquad \text{where } (s,t) \neq (0,0). \tag{5.2}$$

---

[11] The case $\mathrm{U}(1,1)$ was considered above in Subsection 2.18.

[12] A remark for an expert in algebraic geometry: we consider blow up of the plane $\mathbb{C}^2$ at the point $(-n+\alpha, 0)$.

Substituting this to (3.17), we get

$$\ell_{-n+\alpha+\varepsilon s|\varepsilon t} = (-1)^{n(n-1)/2} 2^{-(\sigma+\tau)n} \prod_{k=1}^{n} \Gamma\bigl(-n+\alpha\varepsilon(s+t)+k\bigr)\times$$

$$\times \sum_{\mathbf{m}} \left\{ \frac{(-1)^{\sum m_j} \prod\limits_{1\leqslant a<b\leqslant n} (m_a-m_b)}{\prod\limits_{k=1}^{n} \Gamma(\alpha+\varepsilon s-m_k)\Gamma(\varepsilon t+m_k+1)} \chi_{\mathbf{m}}(g) \right\}. \qquad (5.3)$$

Theorem 5.3 *a) Let $s \neq -t$. Then there exist a limit in the sense of distributions:*

$$\ell^{s:t}(z) := \lim_{\varepsilon\to 0} \ell_{-n+\alpha+\varepsilon s|\varepsilon t}(z), \qquad (5.4)$$

In other words, the function $(\sigma|\tau) \mapsto \ell_{\sigma|\tau}$ has a removable singularity at $\varepsilon = 0$ on the line

$$\sigma = -n+\alpha+\varepsilon s, \qquad \tau = \varepsilon t, \qquad \text{where } \varepsilon \in \mathbb{C}.$$

b) Denote by $c_{\mathbf{m}}(s:t)$ the Fourier coefficients of $\ell^{s:t}$. If $\mathbf{m}$ is in the 'tail', i.e., $\mathbf{m} \notin \cup Z_j$, then $c_{\mathbf{m}}(s:t) = 0$.

c) Moreover, $\ell^{s:t}$ admits a decomposition

$$\ell^{s:t} = \sum_{j=0}^{n-\alpha} \frac{t^j s^{n-\alpha-j}}{(s+t)^{n-\alpha}} \mathfrak{L}_j, \qquad (5.5)$$

where $\mathfrak{L}_j$ is of the form

$$\mathfrak{L}_j = \sum_{\mathbf{m}\in Z_j} a_{\mathbf{m}} \chi_{\mathbf{m}}, \qquad (5.6)$$

where $a_{\mathbf{m}}$ do not depend on s, t.

d) For each j all coefficients $a^j_{\mathbf{m}}$ in (5.6) are either positive or negative.

Proof. For the numerator of (5.3) we have the asymptotic

$$\prod_{k=1}^{n} \Gamma\bigl(-n+\alpha\varepsilon(s+t)+k\bigr) = C\,\varepsilon^{-n+\alpha}(s+t)^{-n+\alpha} + O(\varepsilon^{-n+\alpha+1}), \qquad \varepsilon \to 0.$$

Next, examine factors of the denominator,

$$\Gamma(\alpha+\varepsilon s-m_k)\Gamma(\varepsilon t+m_k+1) \sim \begin{cases} A_1(m_k)(\varepsilon t)^{-1} & \text{if } m_k < 0 \\ A_2(m_k) & \text{if } 0 \leqslant m_k < \alpha \ , \\ A_3(m_k)(\varepsilon s)^{-1} & \text{if } m_k \geqslant \alpha, \end{cases} \qquad \varepsilon \to 0$$

where A_1, A_2, A_3 do not depend on s, t. Therefore, the order of the pole of denominator $\prod_k$ of (5.3) is

number of m_j outside the segment $[0, \alpha-1]$

The minimal possible order of a pole of the denominator is $n - \alpha$. In this case, $c_{\mathbf{m}}$ has a finite nonzero limit, of the form

$$c_{\mathbf{m}}(s:t) = A(\mathbf{m}) \cdot \frac{s^{\text{number of } m_k \geqslant \alpha} \cdot t^{\text{number of } m_k < 0.}}{(s+t)^{n-\alpha}}$$

If an order of pole in the denominator is $> n - \alpha$, then $c_{\mathbf{m}}(s:t) = 0$. This corresponds to the tail.

We omit to watch the positivity of $c_{\mathbf{m}}(s:t)$.

Formally, it is necessary to watch the growth of $c_{\mathbf{m}}(s:t)$ as $\mathbf{m} \to \infty$ and the growth of

$$\frac{\partial}{\partial \varepsilon} c_{\mathbf{m}}(-n + \alpha + \varepsilon s, \varepsilon t)$$

to be sure that (5.4) is a limit in the sense of distributions. This is a more-or-less trivial exercise on the Gamma-function. $\square$

There are many ways to express $\mathfrak{L}^j$ in the terms of $\ell^{s:t}$. One of variants is given in the following obvious proposition.

Proposition 5.4 *The distribution $\mathfrak{L}_j$ is given by the formula*

$$\mathfrak{L}_j(z) = \frac{1}{j!} \frac{\partial^j}{\partial t^j} (1+t)^{n-\alpha} \ell^{1:t}(z) \Big|_{t=0} \tag{5.7}$$

5.3. The family of invariant Hermitian forms. Thus, for $(\sigma, \tau) = (-n + \alpha, 0)$ we obtained the following families of $\rho_{-n+\alpha|0}$-invariant Hermitian forms

$$R^{s:t}(f_1, f_2) := \iint_{\mathrm{U}(n) \times \mathrm{U}(n)} \ell^{s:t}(zu^*) f_1(z) \overline{f_2(u)} \, d\mu(z) \, d\mu(u) \tag{5.8}$$

and

$$Q_j(f_1, f_2) := \iint_{\mathrm{U}(n) \times \mathrm{U}(n)} \mathfrak{L}_j(zu^*) f_1(z) \overline{f_2(u)} \, d\mu(z) \, d\mu(u) \tag{5.9}$$

They are related as

$$R^{s:t}(f_1, f_2) = \sum_{j=0}^{n-\alpha} \frac{t^j s^{n-\alpha-j}}{(s+t)^{n-\alpha}} \mathfrak{L}_j(f_1, f_2)$$

A form $\mathfrak{L}_j$ is zero on

$$Y_j := W_{tail} \oplus (\oplus_{i \neq j} W_i)$$

and determines an inner product on $W_j \simeq C^\infty(\mathrm{U}(n))/Y_j$.

6 Some problems of harmonic analysis

6.1. Tensor products $\rho_{\sigma|\tau} \otimes \rho_{\sigma'|\tau'}$. Nowadays the problem of decomposition of a tensor product of two arbitrary unitary representations does not seem interesting. We propose several informal arguments for reasonableness of the problem in our case.

a) For $n = 1$ it is precisely the well-known problem of decomposition of tensor products of unitary representations of $\mathrm{SL}(2,\mathbb{R})^\sim \simeq \mathrm{SU}(1,1)^\sim$, see [36], [23], [39], [16], [32].

b) Decomposition of tensor products $\rho_{\sigma,0} \otimes \rho_{\sigma',0}$ of holomorphic representations is a well-known combinatorial problem, see [19].

c) Tensor products $\rho_{\sigma,0} \otimes \rho_{0|\tau'}$ are Berezin representations, see [4], [47], [27].

d) All problems a)–c) have interesting links with theory of special functions.

e) There is a canonical isomorphism ${}^{13})$

$$\rho_{-n/2|-n/2} \otimes \rho_{-n/2|-n/2} \simeq L^2\bigl(\mathrm{U}(n,n)/\mathrm{GL}(n,\mathbb{C})\bigr). \tag{6.1}$$

Thus, we again come to a classical problem, i.e., the problem of decomposition of L^2 on a pseudo-Riemannian symmetric space G/H, see [11], [35].[14] General tensor products $\rho_{\sigma|\tau} \otimes \rho_{\sigma'|\tau'}$ can be regarded as deformations of the space $L^2\bigl(\mathrm{U}(n,n)/\mathrm{GL}(n,\mathbb{C})\bigr)$.

6.2. Restriction problems.

1.Consider the group $G^* := \mathrm{U}(n,n)$ and its subgroup $G := \mathrm{O}(n,n)$. The group G has an open dense orbit on the space $\mathrm{U}(n)$, namely

$$G/H := \mathrm{O}(n,n)/\mathrm{O}(n,\mathbb{C}).$$

The restriction of the representation $\rho_{-n/2|-n/2}$ to G is equivalent to the representation of G in $L^2(G/H)$. Restrictions of other $\rho_{\sigma|\tau}$ can be regarded as deformations of $L^2(G/H)$.

The same argument produces deformations of L^2 on some other pseudo-Riemannian symmetric spaces. Precisely, we have the following variants.

2. $G^* = \mathrm{U}(2n,2n)$, $G/H = \mathrm{Sp}(n,n)/\mathrm{Sp}(2n,\mathbb{C})$.

3. $G^* = \mathrm{U}(n,n)$, $G/H = \mathrm{SO}^*(2n)/\mathrm{O}(n,\mathbb{C})$.

4. $G^* = \mathrm{U}(2n,2n)$, $G/H = \mathrm{Sp}(4n,\mathbb{R})/\mathrm{Sp}(2n,\mathbb{C})$.

5. $G^* = \mathrm{U}(p+q,p+q)$, $G/H = \mathrm{U}(p,q) \times \mathrm{U}(p,q)/\mathrm{U}(p,q)$. In this case, $G/H \simeq \mathrm{U}(p,q)$.

6. $G^* = \mathrm{U}(n,n)$, $G = \mathrm{GL}(n,\mathbb{C})$. In this case, we have $(n+1)$ open orbits $G/H_p = \mathrm{GL}(n,\mathbb{C})/\mathrm{U}(p,n-p)$.

---

[13] Indeed, $\mathrm{U}(n) \simeq \mathrm{U}(n,n)/P$, where P is a maximal parabolic subgroup in $\mathrm{U}(n,n)$. The group $\mathrm{U}(n,n)$ has an open orbit on $\mathrm{U}(n,n)/P \times \mathrm{U}(n,n)/P$, the stabilizer of a point is $\simeq \mathrm{GL}(n,\mathbb{C})$.

[14] In a certain sense, the Plancherel formula for $L^2(G/H)$ was obtained in [1], [6]. However no Plancherel measure, nor spectra are known. The corresponding problems remain open.

7. $G^* = \mathrm{U}(n,n)\times \mathrm{U}(n,n)$, $G=\mathrm{U}(n,n)$. This is the problem about tensor products discussed above.

6.3. The Gelfand–Gindikin programm. Recall the statement of the problem, see [13], [34]. Let G/H be a pseudo-Riemannian symmetric space. The natural representation of G in $L^2(G/H)$ has several pieces of spectrum. Therefore, $L^2(G/H)$ admits a natural orthogonal decomposition into direct summands having uniform spectra. The problem is: *to describe explicitly the corresponding subspaces or corresponding projectors.*

In Subsection 5.1 we have obtained a natural decomposition of $L^2(\mathrm{U}(n))$ into $(n+1)$ direct summands. Therefore *in the cases listed in Subsection 6.2 we have a natural orthogonal decompositions of* $L^2(G/H)$.

In any case, for the one-sheet hyperboloid $\mathrm{U}(1,1)/\mathbb{C}^*$ we get the desired construction (see Molchanov [24], [25]).

6.4. Matrix Sobolev spaces? Our inner product $\langle\cdot,\cdot\rangle_{\sigma|\tau}$ seems to be similar to Sobolev-type inner products discussed in Subsection2.13. However it is not a Sobolev inner product, because the kernel $\det(1-zu^*)^{\{\sigma|\tau\}}$ has a non-diagonal singularity.

Denote

$$s=-\sigma-\tau+n.$$

Let F be a distribution on $\mathrm{U}(n)$, let $F=\sum F_{\mathbf{m}}$ be its expansion in a series of elementary harmonics. We have

$$F\in\mathcal{H}_{\sigma|\tau} \qquad \Longleftrightarrow \qquad \sum_{\mathbf{m}} \frac{c_{\mathbf{m}}}{\dim \pi_{\mathbf{m}}} \|F_{\mathbf{m}}\|^2_{L^2}<\infty \qquad \Longleftrightarrow$$

$$\Longleftrightarrow \qquad \sum_{\mathbf{m}} \Bigl\{ \|F_{\mathbf{m}}\|^2_{L^2} \prod_{j=1}^n (1+|m_j|)^s \Bigr\} <\infty, \tag{6.2}$$

where $\|F_{\mathbf{m}}\|_{L^2}$ denotes

$$\|F_{\mathbf{m}}\|_{L^2} := \Bigl(\int_{\mathrm{U}(n)} |F_{\mathbf{m}}(h)|^2\, d\mu(h)\Bigr)^{1/2}$$

Our Hermitian form defines a norm only in the case $|s|<1$, but (6.2) makes sense for arbitrary real s. Thus *we can define a Sobolev space* H_s *on* $\mathrm{U}(n)$ *of arbitrary order.*

Author does not know applications of this remark, but it seems that it can be useful in the following situation.

First, a reasonable harmonic analysis related to semisimple Lie groups is the analysis of unitary representations. But near 1980 Molchanov observed that many identities with special function admit interpretations on "physical level of rigor" as formulas of non-unitary harmonic analysis. Up to now, there are no reasonable interpretations of this phenomenon (but formulas exist, see, e.g. [7], see also [27], Section 1.32 and formula (2.6)–(2.15)). In particular, we do not know reasonable functional spaces that can be scene of action of such analysis. It seems that our spaces H_s can be possible candidates.

References

[1] van den Ban, E.P., Schlichtkrull, H., *The most continuous part of the Plancherel decomposition for a reductive symmetric space.* Ann. Math., 145 (1997), 267–364

[2] Bargmann, V. *Irreducible unitary representations of the Lorentz group.* Ann. Math, 48 (1947), 568–640

[3] Berezin, F.A., *Quantization in complex symmetric spaces.* Izv. Akad. Nauk SSSR, Ser. Math., 39, 2, 1362–1402 (1975); English translation: Math USSR Izv. 9 (1976), No 2, 341–379(1976)

[4] Berezin, F. A. *The connection between covariant and contravariant symbols of operators on classical complex symmetric spaces.* Sov. Math. Dokl. 19 (1978), 786–789

[5] Branson, Th., Olafsson, G., Orsted, B. *Spectrum generating operators and intertwining operators for representations induced from a maximal parabolic subgroup,* J. Funct. Anal., 135, 163–205.

[6] Delorme, P. *Formule de Plancherel pour les espaces symmétrique reductifs.* Ann. Math., 147 (1998), 417–452

[7] van Dijk, G., Molchanov, V.F. *The Berezin form for rank one para-Hermitian symmetric spaces.* J. Math. Pure. Appl., 78 (1999), 99–119.

[8] Dvorsky, A., Sahi, S. *Explicit Hilbert spaces for certain unipotent representations. II.* Invent. Math. 138 (1999), no. 1, 203–224.

[9] Dvorsky, A., Sahi, S. *Explicit Hilbert spaces for certain unipotent representations. III.* J. Funct. Anal. 201(2003), no. 2, 430–456.

[10] Faraut, J., Koranyi, A., *Analysis in symmetric cones.* Oxford Univ.Press, (1994)

[11] Flensted-Jensen, M. *Discrete series for semisimple symmetric spaces.* Ann. of Math. (2) 111 (1980), no. 2, 253–311.

[12] Friedrichs, K. O. *Mathematical aspects of the quantum theory of fields.* Interscience Publishers, Inc., New York, 1953.

[13] Gelfand, I. M., Gindikin, S. G. *Complex manifolds whose skeletons are semisimple Lie groups and analytic discrete series of representations.* Funct. Anal. Appl., 11 (1978), 258–265

[14] Gelfand, I.M., Naimark, M.I., *Unitary representations of classical groups. Unitary representations of classical groups.* Trudy MIAN., t.36 (1950); German translation: Gelfand I.N., Neumark M.A., *Unitare Darstellungen der klassischen gruppen.*, Akademie-Verlag, Berlin, 1957.

[15] Gradshtein, I.S., Ryzhik, I.M. *Tables of integrals, sums and products.* Fizmatgiz, 1963; English translation: Acad. Press, NY, 1965

[16] Groenevelt, W., Koelink, E., Rosengren, H. *Continuous Hahn polynomials and Clebsch–Gordan coefficients.* Preprint `http://arxiv.org/math/abs/0302251`

[17] Harish-Chandra, *Representations of semisimple Lie groups IV,* Amer. J. Math., 743–777 (1955). Reprinted in Harish-Chandra *Collected papers,* v.2.

[18] *Higher transcendental functions*, v.1., McGraw-Hill book company, 1953

[19] Jakobsen, H.P., Vergne, M., *Restrictions and expansions of holomorphic representations.* J. Funct. Anal., 34 (1979), 29–53.

[20] Kadell, K. *The Selberg–Jack symmetric functions.* Adv. Math., 130 (1997), 33-102

[21] Kirillov, A.A. *Elements of representation theory,* Moscow, Nauka, 1972; English transl.: Springer, 1976.

[22] Krattenthaler, C. *Advanced determinant calculus.* The Andrews Festschrift (Maratea, 1998). Sem. Lothar. Combin. 42 (1999), Art. B42q, 67 pp. (electronic).

[23] Molchanov, V. F.*Tensor products of unitary representations of the three-dimensional Lorentz group.* Izv. Akad. Nauk SSSR Ser. Mat. 43 (1979), no. 4, 860–891, 967. English transl. in Izvestia.

[24] Molchanov, V. F. *Quantization on the imaginary Lobachevsky plane.* Funct. Anal. Appl., 14 (1980), 162–144

[25] Neretin, Yu. A. *The restriction of functions holomorphic in a domain to a curve lying in the boundary, and discrete* $\mathrm{SL}_2(\mathbb{R})$*-spectra.* Izvestia: Mathematics, 62:3(1998), 493–513

[26] Neretin, Yu.A., *Matrix analogs of* B*-function and Plancherel formula for Berezin kernel representations,* Mat. Sbornik, 191, No.5 (2000), 67–100;

[27] Neretin, Yu.A., *Plancherel formula for Berezin deformation of* L^2 *on Riemannian symmetric space*, J. Funct. Anal. (2002), 189(2002), 336–408.

[28] Neretin, Yu.A. *Matrix balls, radial analysis of Berezin kernels, and hypergeometric determinants,* Moscow Math. J., v.1 (2001), 157–221.

[29] Neretin, Yu.A. *Notes Sahi–Stein representations and some problems of non-*L^2 *harmonic analysis.*, J. Math. Sci., New York, 141 (2007), 1452–1478

[30] Neretin, Yu. A. *Notes on matrix analogs of Sobolev spaces and Stein–Sahi representations.* Preprint, `http://arxiv.org/abs/math/0411419`

[31] Neretin, Yu. A. *Lectures on Gaussian integral operators and classical groups,* to appear.

[32] Neretin, Yu. A. *Some continuous analogs of expansion in Jacobi polynomials and vector-valued orthogonal bases.* Funct. Anal. Appl., 39 (2005), 31–46.

[33] Neretin, Yu.A., Olshanskii, G.I., *Boundary values of holomorphic functions , singular unitary representations of groups $O(p, q)$ and their limits as $q \to \infty$.* Zapiski nauchn. semin. POMI RAN 223, 9–91(1995); English translation: J.Math.Sci., New York, 87, 6 (1997), 3983–4035.

[34] Olshanskij, G.I., *Complex Lie semigroups, Hardy spaces, and Gelfand–Gindikin programm.* Deff. Geom. Appl., 1 (1991), 235–246

[35] Oshima, T. *A calculation of c-functions for semisimple symmetric spaces. Lie groups and symmetric spaces,* 307–330, Amer. Math. Soc. Transl. Ser. 2, 210, Amer. Math. Soc., Providence, RI, 2003.

[36] Pukanszky, L., *On the Kronecker products of irreducible unitary representations of the* 2×2 *real unimodular group.* Trans. Amer. Math. Soc., 100 (1961), 116–152

[37] Pukanzsky, L. *Plancherel formula for universal covering group of* $\mathrm{SL}(2, \mathbb{R})$. Math. Ann., 156 (1964), 96-?

[38] Ricci, F., Stein, E. M. *Homogeneous distributions on spaces of Hermitean matrices.* J. Reine Angew. Math. 368 (1986), 142–164.

[39] Rosengren, H. *Multilinear Hankel forms of higher order and orthogonal polynomials.* Math. Scand., 82 (1998), 53-88

[40] Sahi, S. *A simple construction of Stein's complementary series representations.* Proc. Amer. Math. Soc. 108 (1990), no. 1, 257–266.

[41] Sahi, S., *Unitary representations on the Shilov boundary of a symmetric tube domain,* Contemp. Math. 145 (1993) 275–286.

[42] Sahi, S. *Jordan algebras and degenerate principal series*, J. Reine Angew.Math. 462 (1995) 1–18.

[43] Sahi, S. *Explicit Hilbert spaces for certain unipotent representations.* Invent. Math. 110 (1992), no. 2, 409–418.

[44] Sahi, S., Stein, E. M. *Analysis in matrix space and Speh's representations.* Invent. Math. 101 (1990), no. 2, 379–393.

[45] Sally, P. J., *Analytic continuations of irreducible unitary representations of the universal covering group of* $\mathrm{SL}(2, \mathbb{R})$. Amer. Math. Soc., Providence, 1967

[46] Stein, E. M. *Analysis in matrix spaces and some new representations of* $\mathrm{SL}(N, C)$. Ann. of Math. (2) 86 1967 461–490.

[47] Unterberger, A., Upmeier, H., *The Berezin transform and invariant differential operators.* Comm.Math.Phys.,164, 563–597(1994)

[48] Vogan, D. A., *The unitary dual of* $\mathrm{GL}(n)$ *over an Archimedean field.* Invent. Math. 83 (1986), no. 3, 449–505.

[49] Weyl, H. *The Classical Groups. Their Invariants and Representations.* Princeton University Press, Princeton, N.J., 1939.

University of Vienna, Math. Dept.,
Nordbergstarsse, 15, Vienna, Austria
& Math. Phys.Group, Institute for the Theoretical and Experimental Physics,
Bolshaya Cheremushkinskaya, 25,
Moscow 117 259, Russia
& TFFA, MechMath Dept., Moscow State University,
Vorob'evy Gory, Moscow, Russia
`neretinmccme.ru`
`URL:www.mat.univie.ac.at/∼neretin,`
`wwwth.itep.ru/∼neretin`